

DETERMINANT FUNCTORS ON TRIANGULATED CATEGORIES

MANUEL BREUNING

ABSTRACT. We study determinant functors which are defined on a triangulated category and take values in a Picard category. The two main results are the existence of a universal determinant functor for every small triangulated category, and a comparison theorem for determinant functors on a triangulated category with a non-degenerate bounded t-structure and determinant functors on its heart. For a small triangulated category $\mathcal{T}$ we give a natural definition of groups $K_0(\mathcal{T})$ and $K_1(\mathcal{T})$ in terms of the universal determinant functor on $\mathcal{T}$, and we show that $K_i(\mathcal{T}) \cong K_i(\mathcal{E})$ for $i = 0$ and 1 if $\mathcal{T}$ has a non-degenerate bounded t-structure with heart $\mathcal{E}$.

1. INTRODUCTION

The systematic study of determinant functors began with the construction of a determinant functor on the category of perfect complexes of $\mathcal{O}_X$ -modules on a scheme X by Knudsen and Mumford [8]. For every perfect complex A they defined a graded invertible $\mathcal{O}_X$ -module $f(A)$ (which is essentially the alternating product of the top exterior powers of a bounded locally free resolution of A), and they showed that f can be made into a functor if one only considers quasi-isomorphisms of perfect complexes. Furthermore for every short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ of perfect complexes they defined an isomorphism $f(B) \cong f(A) \otimes f(C)$ in the category of graded invertible $\mathcal{O}_X$ -modules which satisfies a list of natural properties.

The functor f factors through the category whose objects are perfect complexes of $\mathcal{O}_X$ -modules and whose morphisms are isomorphisms in the derived category of $\mathcal{O}_X$ -modules. However it was observed by Knudsen and Mumford [8, p. 44] that in general it is not possible to define a natural isomorphism $f(B) \cong f(A) \otimes f(C)$ for every distinguished triangle $A \rightarrow B \rightarrow C \rightarrow TA$ of perfect complexes.

The results of Knudsen and Mumford were extended to a theory of determinant functors on exact categories with values in (commutative) Picard categories (see Deligne [5], Knudsen [7]). These determinant functors have found applications in various areas of mathematics. However since one is often interested in applying determinant functors to objects in a derived category, one would like to have determinant functors which are not only well-behaved on short exact sequences but also take into account the triangulated structure of these categories. The purpose of this paper is to suggest a definition of determinant functors on triangulated categories and to demonstrate that this definition leads to an interesting theory.

Recall that a Picard category $\mathcal{P}$ is a category with a bifunctor $\otimes$ and associativity and commutativity isomorphisms satisfying several natural properties. We define a determinant functor f on a triangulated category $\mathcal{T}$ with values in a Picard

Date: October 13, 2006.

The author was supported by EPSRC grant GR/S91772/01.

category $\mathcal{P}$, in symbols $f : \mathcal{T} \rightarrow \mathcal{P}$, to be a pair $f = (f_1, f_2)$ consisting of a functor $f_1 : \mathcal{T}_{\text{iso}} \rightarrow \mathcal{P}$ (where $\mathcal{T}_{\text{iso}}$ is the category obtained from $\mathcal{T}$ by restricting morphisms to isomorphisms) together with an isomorphism $f_2(\Delta) : f_1(B) \cong f_1(A) \otimes f_1(C)$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$, such that f_2 is natural in isomorphisms of distinguished triangles and compatible with direct sums and octahedral diagrams in $\mathcal{T}$. We will also define morphisms of determinant functors and therefore obtain a category $\det(\mathcal{T}, \mathcal{P})$ of determinant functors on $\mathcal{T}$ with values in $\mathcal{P}$.

A determinant functor $f : \mathcal{T} \rightarrow \mathcal{V}$ is called universal if for every Picard category $\mathcal{P}$ the induced functor

$$\text{Hom}^{\otimes}(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$$

which sends M to $M \circ f$ is an equivalence of categories. Here $\text{Hom}^{\otimes}(\mathcal{V}, \mathcal{P})$ denotes the category of monoidal functors from $\mathcal{V}$ to $\mathcal{P}$, and $M \circ f$ is the natural composition of a determinant functor and a monoidal functor. Every small triangulated category $\mathcal{T}$ has a universal determinant functor $f : \mathcal{T} \rightarrow \mathcal{V}$. We will prove this by giving an explicit construction of $\mathcal{V}$ in terms of generators and relations.

The existence of a universal determinant functor $f : \mathcal{E} \rightarrow \mathcal{V}$ on a small exact category $\mathcal{E}$ was shown by Deligne [5, §4]. His proof shows that for $i = 0$ and 1 the Quillen K -group $K_i(\mathcal{E})$ is canonically isomorphic to the homotopy group $\pi_i(\mathcal{V})$. For a small triangulated category $\mathcal{T}$ this motivates the definition $K_i(\mathcal{T}) = \pi_i(\mathcal{V})$ for $i = 0, 1$ where $f : \mathcal{T} \rightarrow \mathcal{V}$ is any universal determinant functor on $\mathcal{T}$. It follows easily from the definition of a universal determinant functor that these K -groups are well-defined and functorial. Furthermore it is not difficult to see that $K_0(\mathcal{T})$ is the Grothendieck group of $\mathcal{T}$.

Next we assume that $\mathcal{T}$ is a triangulated category with a non-degenerate bounded t -structure and we let $\mathcal{E}$ be its heart. We will show that for every Picard category $\mathcal{P}$ there is a natural equivalence of categories

$$\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P}).$$

If in addition $\mathcal{T}$ is small we deduce that $K_i(\mathcal{T}) \cong K_i(\mathcal{E})$ for $i = 0, 1$. The proof that the functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ is an equivalence of categories contains several intermediate steps of independent interest. For example we will show that any bounded cohomological functor on a triangulated category induces a functor between certain categories of determinant functors.

We should mention two topics which are not addressed in this paper. Firstly we do not give any applications of determinant functors on triangulated categories. One application which will be discussed elsewhere is a new construction of certain Euler characteristics in relative algebraic K_0 -groups (which are defined in [4], [3] and [2]) using universal determinant functors on triangulated categories of perfect complexes. Secondly many basic questions regarding the groups K_0 and K_1 of a triangulated category remain open. In particular we do not know how the group K_1 is related to the K_1 -groups of a triangulated category defined by Neeman [10].

Finally we would like to remark that results in the spirit of this paper have also been shown for Waldhausen categories instead of triangulated categories. Further details on this can be found in the paper by Muro and Tonks [9] and in forthcoming work by Witte.

This paper is organized as follows. In §2 we recall the definitions of Picard categories, monoidal functors and determinant functors on exact categories, and

we fix our notation for triangulated categories. The definition and basic properties of determinant functors on triangulated categories are given in §3. Next in §4 we show the existence of a universal determinant functor for every small triangulated category and we define the groups $K_i(\mathcal{T})$ for $i = 0$ and 1 . Finally in §5 we prove the comparison result for determinant functors on a triangulated category with a non-degenerate bounded t-structure and determinant functors on its heart.

I would like to thank David Burns and Matthias Flach for helpful discussions.

2. PRELIMINARIES

In this section we collect some necessary background material. We first summarize the relevant facts about Picard categories and monoidal functors between Picard categories. We then recall the definition of determinant functors on exact categories. Finally we fix the terminology that we use for triangulated categories.

2.1. Picard categories. An AC tensor category is a category $\mathcal{P}$ together with a bifunctor $\otimes : \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$ and compatible associativity and commutativity constraints φ and ψ respectively, i.e. $\varphi_{X,Y,Z} : X \otimes (Y \otimes Z) \xrightarrow{\cong} (X \otimes Y) \otimes Z$ and $\psi_{X,Y} : X \otimes Y \xrightarrow{\cong} Y \otimes X$ are natural isomorphisms satisfying $\psi_{Y,X} \circ \psi_{X,Y} = \text{id}_{X \otimes Y}$, the pentagonal axiom [11, Chap. I, 1.1.1] and the hexagonal axiom [11, Chap. I, 2.1.1]. We will write $\mathcal{P}$ for an AC tensor category $(\mathcal{P}, \otimes, \varphi, \psi)$ if $\otimes, \varphi$ and ψ are clear from the context. The coherence theorem for AC tensor categories often allows us to omit explicit reference to φ and ψ and to write expressions involving $\otimes$ without parentheses; see [11, Chap. I, 2.1.2] for a precise formulation.

Definition 2.1. An AC tensor category $\mathcal{P} = (\mathcal{P}, \otimes, \varphi, \psi)$ is called a Picard category if it satisfies the following three conditions: $\mathcal{P}$ is non-empty, every morphism in $\mathcal{P}$ is an isomorphism, and for every object W in $\mathcal{P}$ the functor $X \mapsto W \otimes X$ is an autoequivalence of $\mathcal{P}$.

We remark that such a category is sometimes called a commutative Picard category. In this paper we will not consider any non-commutative Picard categories and we will therefore always omit the word commutative.

A unit in a Picard category $\mathcal{P}$ is a triple (U, λ, ρ) consisting of an object U and natural isomorphisms $\lambda_X : X \xrightarrow{\cong} U \otimes X$ and $\rho_X : X \xrightarrow{\cong} X \otimes U$ satisfying the conditions [11, Chap. I, 1.3.1 and 2.2.1] (where the notation g and d is used instead of λ and ρ). A morphism of units $(U, \lambda, \rho) \rightarrow (U', \lambda', \rho')$ is a morphism $\alpha : U \rightarrow U'$ such that $(\alpha \otimes \text{id}_X) \circ \lambda_X = \lambda'_X$ and $(\text{id}_X \otimes \alpha) \circ \rho_X = \rho'_X$ for all objects X . A reduced unit in $\mathcal{P}$ is a pair (U, δ) where U is an object and $\delta : U \xrightarrow{\cong} U \otimes U$ is an isomorphism. A morphism of reduced units $(U, \delta) \rightarrow (U', \delta')$ is a morphism $\alpha : U \rightarrow U'$ such that $(\alpha \otimes \alpha) \circ \delta = \delta' \circ \alpha$. By [11, Chap. I, 2.2.5] the functor from the category of units to the category of reduced units which sends (U, λ, ρ) to (U, λ_U) (note that $\lambda_U = \rho_U$) is an isomorphism of categories. We will usually identify a reduced unit (U, δ) with the unique unit (U, λ, ρ) for which $\delta = \lambda_U = \rho_U$, and in particular call the pair (U, δ) a unit. To simplify the notation we will sometimes write δ for the isomorphisms $\lambda_X : X \rightarrow U \otimes X$ and $\rho_X : X \rightarrow X \otimes U$.

There exists a natural product of two units. If (U, λ, ρ) and (U', λ', ρ') are two units then we define their product $(U, \lambda, \rho) \otimes (U', \lambda', \rho')$ to be the unit $(U \otimes U', \lambda'', \rho'')$ where λ''_X is the composite $X \xrightarrow{\lambda'_X} U' \otimes X \xrightarrow{\lambda_U \otimes \text{id}} (U \otimes U') \otimes X$ and ρ''_X is the

composite $X \xrightarrow{\rho_X} X \otimes U \xrightarrow{\text{id} \otimes \rho'_U} X \otimes (U \otimes U')$. In terms of reduced units the product has the following description. If (U, δ) and (U', δ') are two (reduced) units in $\mathcal{P}$ then their product is $(U, \delta) \otimes (U', \delta') = (U \otimes U', \delta'')$ where δ'' is the composite isomorphism $U \otimes U' \xrightarrow{\delta \otimes \delta'} (U \otimes U) \otimes (U' \otimes U') \cong (U \otimes U') \otimes (U \otimes U')$ (the last isomorphism does not depend on the order of the two copies of U and of U' because $\psi_{U,U} = \text{id}_{U \otimes U}$ and $\psi_{U',U'} = \text{id}_{U' \otimes U'}$).

Every Picard category has a unit and units are unique up to unique isomorphism. Therefore computations with units can also be performed by fixing a unit in $\mathcal{P}$. If $(1_{\mathcal{P}}, \delta)$ with $\delta : 1_{\mathcal{P}} \rightarrow 1_{\mathcal{P}} \otimes 1_{\mathcal{P}}$ is a fixed unit in $\mathcal{P}$ then the category of units is isomorphic to the category of pairs (U, ω) where U is an object in $\mathcal{P}$ and $\omega : U \rightarrow 1_{\mathcal{P}}$ is an isomorphism, and where a morphism $(U, \omega) \rightarrow (U', \omega')$ is a morphism $\alpha : U \rightarrow U'$ such that $\omega' \circ \alpha = \omega$. The product of (U, ω) and (U', ω') is given by $(U \otimes U', \omega'')$ where ω'' is the composite $U \otimes U' \xrightarrow{\omega \otimes \omega'} 1_{\mathcal{P}} \otimes 1_{\mathcal{P}} \xrightarrow{\delta^{-1}} 1_{\mathcal{P}}$. Since in general a Picard category does not have a unique (or a canonical) unit, we prefer not to work with a fixed unit in this paper.

The homotopy groups $\pi_0(\mathcal{P})$ and $\pi_1(\mathcal{P})$ of a small Picard category $\mathcal{P}$ are defined as follows: $\pi_0(\mathcal{P})$ is the group of isomorphism classes of objects of $\mathcal{P}$ (with product induced by $\otimes$) and $\pi_1(\mathcal{P})$ is the group of automorphisms of any object X of $\mathcal{P}$ (this is well-defined because for any two objects X and Y there exists a canonical isomorphism $\text{Aut}_{\mathcal{P}}(X) \cong \text{Aut}_{\mathcal{P}}(Y)$). The groups $\pi_0(\mathcal{P})$ and $\pi_1(\mathcal{P})$ are abelian. We define a map $\varepsilon : \pi_0(\mathcal{P}) \rightarrow \pi_1(\mathcal{P})$ by sending the isomorphism class of an object X to the automorphism $\psi_{X,X} \in \text{Aut}_{\mathcal{P}}(X \otimes X) = \pi_1(\mathcal{P})$. It is not difficult to show that ε is a homomorphism.

2.2. Monoidal functors. A monoidal functor from a Picard category $(\mathcal{P}, \otimes, \varphi, \psi)$ to a Picard category $(\mathcal{P}', \otimes', \varphi', \psi')$ is a pair (M, c) consisting of a functor $M : \mathcal{P} \rightarrow \mathcal{P}'$ and a natural isomorphism $c_{X,Y} : M(X) \otimes' M(Y) \xrightarrow{\cong} M(X \otimes Y)$ which is compatible with the associativity and commutativity constraints (see [11, Chap. I, 4.2.1, 4.2.2]). We shall write M for the monoidal functor (M, c) if no confusion can occur. If $(M, c), (N, d)$ are monoidal functors from $\mathcal{P}$ to $\mathcal{P}'$ then a morphism from (M, c) to (N, d) is a natural transformation $\kappa : M \rightarrow N$ which satisfies $\kappa_{X \otimes Y} \circ c_{X,Y} = d_{X,Y} \circ (\kappa_X \otimes' \kappa_Y)$. Every morphism of monoidal functors of Picard categories is automatically an isomorphism. We write $\text{Hom}^{\otimes}(\mathcal{P}, \mathcal{P}')$ for the category whose objects are monoidal functors from $\mathcal{P}$ to $\mathcal{P}'$ and whose morphisms are morphisms of monoidal functors. Finally we note that if $(M, c) : \mathcal{P} \rightarrow \mathcal{P}'$ and $(M', c') : \mathcal{P}' \rightarrow \mathcal{P}''$ are monoidal functors then there is an obvious way to define the composition $(M', c') \circ (M, c) : \mathcal{P} \rightarrow \mathcal{P}''$.

A monoidal functor $(M, c) : \mathcal{P} \rightarrow \mathcal{P}'$ of small Picard categories induces homomorphisms $\pi_i(M) : \pi_i(\mathcal{P}) \rightarrow \pi_i(\mathcal{P}')$ for $i = 0, 1$. It is not difficult to see that two isomorphic monoidal functors induce the same homomorphisms on π_i for $i = 0, 1$.

Lemma 2.2. *For a monoidal functor $(M, c) : \mathcal{P} \rightarrow \mathcal{P}'$ of small Picard categories the following are equivalent.*

- (i) *M is an equivalence of categories, i.e. there exists a functor $N : \mathcal{P}' \rightarrow \mathcal{P}$ (not a monoidal functor) such that $N \circ M$ and $M \circ N$ are isomorphic to the respective identity functors.*
- (ii) *(M, c) is an equivalence of Picard categories, i.e. there exists a monoidal functor $(N, d) : \mathcal{P}' \rightarrow \mathcal{P}$ such that $(N, d) \circ (M, c)$ and $(M, c) \circ (N, d)$ are*

isomorphic to the respective identity functors (isomorphic as monoidal functors).

(iii) (M, c) induces isomorphisms on π_0 and π_1 .

Proof. The implication (ii) $\implies$ (i) is obvious, and (i) $\implies$ (iii) holds because if $M : \mathcal{P} \rightarrow \mathcal{P}'$ is an equivalence of categories then M induces a bijection on the set of isomorphism classes of $\mathcal{P}$ and $\mathcal{P}'$, and an isomorphism $\text{Aut}_{\mathcal{P}}(X) \cong \text{Aut}_{\mathcal{P}'}(M(X))$ for any object X of $\mathcal{P}$.

To show (iii) $\implies$ (ii) we construct a monoidal functor $(N, d) : \mathcal{P}' \rightarrow \mathcal{P}$ as follows. For every object X' of $\mathcal{P}'$ we fix an object $N(X')$ in $\mathcal{P}$ and an isomorphism $\kappa_{X'} : M(N(X')) \rightarrow X'$ in $\mathcal{P}'$ (such $N(X')$ and $\kappa_{X'}$ exist because M induces a surjection on isomorphism classes). Since M induces an injection on isomorphism classes and a bijection on automorphisms, it follows that for any isomorphism $\alpha' : X' \rightarrow Y'$ in $\mathcal{P}'$ there exists a unique isomorphism $N(\alpha') : N(X') \rightarrow N(Y')$ such that $\alpha' \circ \kappa_{X'} = \kappa_{Y'} \circ M(N(\alpha'))$. Furthermore for every pair X', Y' there exists a unique $d_{X', Y'} : N(X') \otimes N(Y') \rightarrow N(X' \otimes Y')$ such that $\kappa_{X'} \otimes \kappa_{Y'} = \kappa_{X' \otimes Y'} \circ M(d_{X', Y'}) \circ c_{N(X'), N(Y')}$. One easily verifies that $(N, d) : \mathcal{P}' \rightarrow \mathcal{P}$ is a monoidal functor and that $\kappa : (M, c) \circ (N, d) \rightarrow \text{id}_{\mathcal{P}'}$ is an isomorphism of monoidal functors. Moreover it is not difficult to see that there also exists an isomorphism $(N, d) \circ (M, c) \cong \text{id}_{\mathcal{P}}$. $\square$

The following lemma describes the automorphisms of a monoidal functor in terms of homomorphisms between certain homotopy groups.

Lemma 2.3. *Let $\mathcal{P}$ and $\mathcal{P}'$ be small Picard categories and $(M, c) : \mathcal{P} \rightarrow \mathcal{P}'$ a monoidal functor. Then there exists a canonical isomorphism*

$$\text{Aut}_{\text{Hom}^{\otimes}(\mathcal{P}, \mathcal{P}')}((M, c)) \cong \text{Hom}(\pi_0(\mathcal{P}), \pi_1(\mathcal{P}'))$$

(where the last Hom is homomorphisms of abelian groups).

Proof. Let $\kappa \in \text{Aut}_{\text{Hom}^{\otimes}(\mathcal{P}, \mathcal{P}')}((M, c))$. Then for every object X in $\mathcal{P}$ we have an automorphism $\kappa_X \in \text{Aut}_{\mathcal{P}'}(M(X)) = \pi_1(\mathcal{P}')$. We define a homomorphism $\varphi : \pi_0(\mathcal{P}) \rightarrow \pi_1(\mathcal{P}')$ by $\varphi([X]) = \kappa_X$, where $[X]$ denotes the class of X in $\pi_0(\mathcal{P})$. This is well-defined as a map of sets because if X and Y are isomorphic in $\mathcal{P}$ then $\kappa_X = \kappa_Y$ in $\pi_1(\mathcal{P}')$ by the naturality of κ . Furthermore φ is a homomorphism because the commutative diagram

$$\begin{array}{ccc} M(X) \otimes' M(Y) & \xrightarrow{c_{X,Y}} & M(X \otimes Y) \\ \downarrow \kappa_X \otimes' \kappa_Y & & \downarrow \kappa_{X \otimes Y} \\ M(X) \otimes' M(Y) & \xrightarrow{c_{X,Y}} & M(X \otimes Y) \end{array}$$

implies that $\kappa_{X \otimes Y} = \kappa_X \cdot \kappa_Y$ in $\pi_1(\mathcal{P}')$.

Conversely, if $\varphi : \pi_0(\mathcal{P}) \rightarrow \pi_1(\mathcal{P}')$ is a homomorphism then setting $\kappa_X = \varphi([X]) \in \pi_1(\mathcal{P}') = \text{Aut}_{\mathcal{P}'}(M(X))$ for an object X in $\mathcal{P}$ defines an automorphism κ of (M, c) . Obviously $\kappa \mapsto \varphi$ and $\varphi \mapsto \kappa$ are mutually inverse isomorphisms. $\square$

2.3. Determinant functors on exact categories. Let $\mathcal{E}$ be an exact category. We will write the exact sequences in $\mathcal{E}$ either as $A \rightarrow B \rightarrow C$ or as $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ and we will always refer to them as short exact sequences. Let w be a class of morphisms in $\mathcal{E}$ which contains all isomorphisms and is closed under composition. The subcategory of $\mathcal{E}$ where morphisms are restricted to w will be denoted by $\mathcal{E}_w$.

Definition 2.4. A determinant functor f on $(\mathcal{E}, w)$ with values in a Picard category $\mathcal{P}$ is a pair $f = (f_1, f_2)$ consisting of a functor $f_1 : \mathcal{E}_w \rightarrow \mathcal{P}$ and for every short exact sequence $\Delta : 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in $\mathcal{E}$ an isomorphism $f_2(\Delta) : f_1(B) \rightarrow f_1(A) \otimes f_1(C)$, such that the following three axioms are satisfied.

(i) *Naturality.* For every morphism of short exact sequences

$$\begin{array}{ccccccc} \Delta : 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \\ & & \downarrow a & & \downarrow b & & \downarrow c \\ \Delta' : 0 & \longrightarrow & A' & \longrightarrow & B' & \longrightarrow & C' \longrightarrow 0 \end{array}$$

with $a, b, c \in w$, the following diagram is commutative.

$$\begin{array}{ccc} f_1(B) & \xrightarrow{f_2(\Delta)} & f_1(A) \otimes f_1(C) \\ \downarrow f_1(b) & & \downarrow f_1(a) \otimes f_1(c) \\ f_1(B') & \xrightarrow{f_2(\Delta')} & f_1(A') \otimes f_1(C') \end{array}$$

(ii) *Associativity.* For every commutative diagram of short exact sequences

$$(1) \quad \begin{array}{ccccc} A & \longrightarrow & B & \longrightarrow & C' \\ \parallel & & \downarrow & & \downarrow \\ A & \longrightarrow & C & \longrightarrow & B' \\ & & \downarrow & & \downarrow \\ & & A' & \xlongequal{\quad} & A' \end{array}$$

the following diagram is commutative

$$\begin{array}{ccc} f_1(C) & \xrightarrow{f_2(\Delta_{h2})} & f_1(A) \otimes f_1(B') \\ \downarrow f_2(\Delta_{v1}) & & \downarrow \text{id} \otimes f_2(\Delta_{v2}) \\ & & f_1(A) \otimes (f_1(C') \otimes f_1(A')) \\ & & \downarrow \varphi_{f_1(A), f_1(C'), f_1(A')} \\ f_1(B) \otimes f_1(A') & \xrightarrow{f_2(\Delta_{h1}) \otimes \text{id}} & (f_1(A) \otimes f_1(C')) \otimes f_1(A'), \end{array}$$

where Δ_{h1} and Δ_{h2} (resp. Δ_{v1} and Δ_{v2}) are the first and second horizontal (resp. vertical) short exact sequences in diagram (1).

(iii) *Commutativity.* For every pair of short exact sequences

$$\begin{aligned} \Delta_1 : A &\rightarrow A \oplus B \rightarrow B, \\ \Delta_2 : B &\rightarrow A \oplus B \rightarrow A \end{aligned}$$

where $A \oplus B$ is a direct sum of A and B and the maps are the canonical inclusions and projections, the following diagram is commutative.

$$\begin{array}{ccc}
 & f_1(A \oplus B) & \\
 f_2(\Delta_1) \swarrow & & \searrow f_2(\Delta_2) \\
 f_1(A) \otimes f_1(B) & \xrightarrow{\psi_{f_1(A), f_1(B)}} & f_1(B) \otimes f_1(A)
 \end{array}$$

Remark 2.5. Our definition of a determinant functor agrees with the definition in [7] because the compatibility axiom in [7, Definition 1.4] follows from the associativity axiom. In [7], w is required to be an SQ-class of morphisms, however the definition makes sense for any class of morphisms w as considered above.

Definition 2.6. Let $f = (f_1, f_2)$ and $g = (g_1, g_2)$ be determinant functors on $(\mathcal{E}, w)$ with values in $\mathcal{P}$. A morphism $f \rightarrow g$ is a natural isomorphism $\lambda : f_1 \rightarrow g_1$ such that $(\lambda_A \otimes \lambda_C) \circ f_2(\Delta) = g_2(\Delta) \circ \lambda_B : f_1(B) \rightarrow g_1(A) \otimes g_1(C)$ for every short exact sequence $\Delta : 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in $\mathcal{E}$.

We will often use the notation $f : (\mathcal{E}, w) \rightarrow \mathcal{P}$ to denote a determinant functor f on $(\mathcal{E}, w)$ with values in $\mathcal{P}$. We write $\det((\mathcal{E}, w), \mathcal{P})$ for the category whose objects are the determinant functors $(\mathcal{E}, w) \rightarrow \mathcal{P}$ and whose morphisms are the morphisms of determinant functors. If $F : (\mathcal{E}', w') \rightarrow (\mathcal{E}, w)$ is an exact functor (i.e. $F : \mathcal{E}' \rightarrow \mathcal{E}$ is an exact functor of exact categories which satisfies $F(w') \subseteq w$) and $M : \mathcal{P} \rightarrow \mathcal{P}'$ is a monoidal functor then we obtain an induced functor $\det((\mathcal{E}, w), \mathcal{P}) \rightarrow \det((\mathcal{E}', w'), \mathcal{P}')$, which sends a determinant functor $f : (\mathcal{E}, w) \rightarrow \mathcal{P}$ to the composite determinant functor $M \circ f \circ F : (\mathcal{E}', w') \rightarrow \mathcal{P}'$ that is defined in the obvious way.

If $w = \text{iso}$ is the class of all isomorphisms in $\mathcal{E}$ then we will in general omit w from the notation. So we write $\det(\mathcal{E}, \mathcal{P})$ instead of $\det((\mathcal{E}, \text{iso}), \mathcal{P})$ and talk about determinant functors on $\mathcal{E}$ with values in $\mathcal{P}$ instead of determinant functors on $(\mathcal{E}, \text{iso})$ with values in $\mathcal{P}$.

2.4. Preliminaries about triangulated categories. Let $\mathcal{T}$ be a triangulated category, i.e. $\mathcal{T}$ is an additive category with an additive automorphism T (the translation functor) and a class of distinguished triangles $A \rightarrow B \rightarrow C \rightarrow TA$ such that the axioms (TR1)–(TR4) in [12] hold.

Let A and B be objects in $\mathcal{T}$. It follows from the axioms of a triangulated category that

$$(2) \quad A \rightarrow A \oplus B \rightarrow B \xrightarrow{0} TA$$

is a distinguished triangle in $\mathcal{T}$, where $A \oplus B$ is a direct sum of A and B , and the maps $A \rightarrow A \oplus B$ and $A \oplus B \rightarrow B$ are the canonical inclusion and projection respectively.

By an octahedral diagram in $\mathcal{T}$ we mean a diagram of the form

$$(3) \quad \begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C' & \longrightarrow & TA \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ A & \longrightarrow & C & \longrightarrow & B' & \xrightarrow{c} & TA \\ & & \downarrow & & \downarrow d & & \\ & & A' & \xlongequal{\quad} & A' & & \\ & & \downarrow e & & \downarrow & & \\ & & TB & \xrightarrow{Tb} & TC' & & \end{array}$$

such that

- (i) the diagram is commutative,
- (ii) the two horizontal triangles and the two vertical triangles are distinguished,
- (iii) the two composite morphisms $B' \xrightarrow{c} TA \xrightarrow{Ta} TB$ and $B' \xrightarrow{d} A' \xrightarrow{e} TB$ are equal.

We remark that octahedral diagrams are the diagrams which appear in the octahedral axiom (TR4) in [12].

An exact functor $(F, \tau) : \mathcal{S} \rightarrow \mathcal{T}$ of triangulated categories is a pair consisting of an additive functor $F : \mathcal{S} \rightarrow \mathcal{T}$ and a natural isomorphism $\tau : F \circ T_{\mathcal{S}} \rightarrow T_{\mathcal{T}} \circ F$ such that if $A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} T_{\mathcal{S}}A$ is a distinguished triangle in $\mathcal{S}$ then $FA \xrightarrow{Fa} FB \xrightarrow{Fb} FC \xrightarrow{\tau_A \circ Fc} T_{\mathcal{T}}FA$ is a distinguished triangle in $\mathcal{T}$. An exact functor sends a distinguished triangle of the form (2) in $\mathcal{S}$ to a distinguished triangle of the same form in $\mathcal{T}$. Furthermore it sends an octahedral diagram in $\mathcal{S}$ to an octahedral diagram in $\mathcal{T}$. For the definition of a morphism of exact functors we refer to [12, Chap. 2, §2, no. 1].

3. DEFINITION AND BASIC PROPERTIES

In this section we define determinant functors on triangulated categories and prove various basic properties for them.

3.1. Definition of determinant functors. Let $\mathcal{T}$ be a triangulated category. We write iso for the class of isomorphisms in $\mathcal{T}$ and $\mathcal{T}_{\text{iso}}$ for the subcategory of $\mathcal{T}$ where morphisms are restricted to isomorphisms.

Definition 3.1. A determinant functor f on $\mathcal{T}$ with values in a Picard category $\mathcal{P}$ is a pair $f = (f_1, f_2)$ consisting of a functor $f_1 : \mathcal{T}_{\text{iso}} \rightarrow \mathcal{P}$ and for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$ an isomorphism $f_2(\Delta) : f_1(B) \rightarrow f_1(A) \otimes f_1(C)$, such that the following three axioms are satisfied.

- (i) *Naturality.* For every isomorphism of distinguished triangles

$$(4) \quad \begin{array}{ccccccc} \Delta : A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & TA \\ \downarrow a & & \downarrow b & & \downarrow c & & \downarrow Ta \\ \Delta' : A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & TA', \end{array}$$

the following diagram is commutative.

$$\begin{array}{ccc} f_1(B) & \xrightarrow{f_2(\Delta)} & f_1(A) \otimes f_1(C) \\ \downarrow f_1(b) & & \downarrow f_1(a) \otimes f_1(c) \\ f_1(B') & \xrightarrow{f_2(\Delta')} & f_1(A') \otimes f_1(C') \end{array}$$

- (ii) *Associativity.* For every octahedral diagram as in (3), the following diagram is commutative

$$\begin{array}{ccc} f_1(C) & \xrightarrow{f_2(\Delta_{h2})} & f_1(A) \otimes f_1(B') \\ \downarrow f_2(\Delta_{v1}) & & \downarrow \text{id} \otimes f_2(\Delta_{v2}) \\ f_1(B) \otimes f_1(A') & \xrightarrow{f_2(\Delta_{h1}) \otimes \text{id}} & f_1(A) \otimes (f_1(C') \otimes f_1(A')) \\ & & \downarrow \varphi_{f_1(A), f_1(C'), f_1(A')} \\ & & (f_1(A) \otimes f_1(C')) \otimes f_1(A') \end{array}$$

where Δ_{h1} and Δ_{h2} (resp. Δ_{v1} and Δ_{v2}) are the first and second horizontal (resp. vertical) distinguished triangles in diagram (3).

- (iii) *Commutativity.* For every pair of distinguished triangles

$$(5) \quad \begin{aligned} \Delta_1 : A &\rightarrow A \oplus B \rightarrow B \xrightarrow{0} TA, \\ \Delta_2 : B &\rightarrow A \oplus B \rightarrow A \xrightarrow{0} TB \end{aligned}$$

where $A \oplus B$ is a direct sum of A and B and the maps are the canonical inclusions and projections, the following diagram is commutative.

$$\begin{array}{ccc} & f_1(A \oplus B) & \\ f_2(\Delta_1) \swarrow & & \searrow f_2(\Delta_2) \\ f_1(A) \otimes f_1(B) & \xrightarrow{\psi_{f_1(A), f_1(B)}} & f_1(B) \otimes f_1(A) \end{array}$$

Definition 3.2. Let $f = (f_1, f_2)$ and $g = (g_1, g_2)$ be determinant functors on $\mathcal{T}$ with values in $\mathcal{P}$. A morphism $f \rightarrow g$ is a natural isomorphism $\lambda : f_1 \rightarrow g_1$ such that $(\lambda_A \otimes \lambda_C) \circ f_2(\Delta) = g_2(\Delta) \circ \lambda_B : f_1(B) \rightarrow g_1(A) \otimes g_1(C)$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$.

We will often use the notation $f : \mathcal{T} \rightarrow \mathcal{P}$ to denote a determinant functor f on $\mathcal{T}$ with values in $\mathcal{P}$. We write $\det(\mathcal{T}, \mathcal{P})$ for the category whose objects are the determinant functors $\mathcal{T} \rightarrow \mathcal{P}$ and whose morphisms are the morphisms of determinant functors. If $F : \mathcal{T}' \rightarrow \mathcal{T}$ is an exact functor of triangulated categories and $M : \mathcal{P} \rightarrow \mathcal{P}'$ is a monoidal functor then we obtain an induced functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{T}', \mathcal{P}')$, which sends a determinant functor $f : \mathcal{T} \rightarrow \mathcal{P}$ to the composite determinant functor $M \circ f \circ F : \mathcal{T}' \rightarrow \mathcal{P}'$ that is defined in the obvious way.

To simplify the notation we will often write $[\cdot]$ for a determinant functor $f = (f_1, f_2)$, i.e. if A (resp. a) is an object (resp. isomorphism) in $\mathcal{T}$ then $[A] = f_1(A)$ (resp. $[a] = f_1(a)$), and if Δ is a distinguished triangle in $\mathcal{T}$ then $[\Delta] = f_2(\Delta)$. This will not cause any confusion.

3.2. Basic properties of determinant functors. Let $[\cdot]$ be a determinant functor on a triangulated category $\mathcal{T}$ with values in a Picard category $\mathcal{P}$.

Lemma 3.3. *If 0 is a zero object in $\mathcal{T}$ then $[0]$ has canonically the structure of a unit in $\mathcal{P}$, i.e. there exists a canonical isomorphism $\delta_0 : [0] \rightarrow [0] \otimes [0]$. If $a : 0 \rightarrow 0'$ is the unique isomorphism of zero objects 0 and $0'$ then $[a] : [0] \rightarrow [0']$ is an isomorphism of units $([0], \delta_0) \rightarrow ([0'], \delta_{0'})$.*

Proof. Applying $[\cdot]$ to the distinguished triangle $0 \rightarrow 0 \rightarrow 0 \rightarrow T0$ gives a canonical isomorphism $\delta_0 : [0] \rightarrow [0] \otimes [0]$ and therefore the structure of a unit on $[0]$. The isomorphism $a : 0 \rightarrow 0'$ induces an isomorphism from the distinguished triangle $0 \rightarrow 0 \rightarrow 0 \rightarrow T0$ to $0' \rightarrow 0' \rightarrow 0' \rightarrow T0'$, and therefore $[a]$ commutes with the unit structures on $[0]$ and $[0']$. $\square$

We remark that the unit structure δ_0 on $[0]$ depends on the zero object 0 in $\mathcal{T}$ and not only on the object $[0]$ in $\mathcal{P}$. Nevertheless to simplify the notation we will talk about the unit $[0]$ when we mean the unit $([0], \delta_0)$.

Lemma 3.4. *Let $a : A \rightarrow B$ be an isomorphism and 0 a zero object in $\mathcal{T}$. Let $\lambda_X : X \rightarrow [0] \otimes X$ and $\rho_X : X \rightarrow X \otimes [0]$ be the natural isomorphisms associated to the unit $[0]$.*

- (i) *Let Δ be the distinguished triangle $A \xrightarrow{a} B \rightarrow 0 \rightarrow TA$. Then $[a] = [\Delta]^{-1} \circ \rho_{[A]} : [A] \rightarrow [B]$.*
- (ii) *Let Δ be the distinguished triangle $0 \rightarrow A \xrightarrow{a} B \rightarrow T0$. Then $[a] = \lambda_{[B]}^{-1} \circ [\Delta] : [A] \rightarrow [B]$.*

Proof. We prove part (ii), the proof of part (i) is similar and left to the reader. Recall that by definition of the unit structure on $[0]$ the isomorphism $\delta_0 = \lambda_{[0]} = \rho_{[0]} : [0] \rightarrow [0] \otimes [0]$ is induced by the distinguished triangle $0 \rightarrow 0 \rightarrow 0 \rightarrow T0$. Now consider the distinguished triangle $\Delta' : 0 \rightarrow B \xrightarrow{\text{id}} B \rightarrow T0$. From the octahedral diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & T0 \\
 \parallel & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & B & \xrightarrow{\text{id}} & B & \longrightarrow & T0 \\
 & & \downarrow \text{id} & & \downarrow \text{id} & & \\
 & & B & \xlongequal{\quad} & B & & \\
 & & \downarrow & & \downarrow & & \\
 & & T0 & \longrightarrow & T0 & &
 \end{array}$$

we obtain $(\delta_0 \otimes \text{id}_{[B]}) \circ [\Delta'] = \varphi_{[0],[0],[B]} \circ (\text{id}_{[0]} \otimes [\Delta']) \circ [\Delta']$. On the other hand by the axioms for a unit $\delta_0 \otimes \text{id}_{[B]} = \varphi_{[0],[0],[B]} \circ (\text{id}_{[0]} \otimes \lambda_{[B]})$. The last two equations imply $[\Delta'] = \lambda_{[B]}$. The isomorphism of distinguished triangles

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A & \xrightarrow{a} & B & \longrightarrow & T0 \\
 \downarrow & & \downarrow a & & \downarrow \text{id} & & \downarrow \\
 0 & \longrightarrow & B & \xrightarrow{\text{id}} & B & \longrightarrow & T0
 \end{array}$$

shows that $[\Delta'] \circ [a] = [\Delta]$. Therefore $[a] = \lambda_{[B]}^{-1} \circ [\Delta]$ as required. $\square$

Lemma 3.5. *Let A be an object of $\mathcal{T}$ and let $-\text{id}_A : A \rightarrow A$ be the negative of the identity on A . Then $[-\text{id}_A] = \varepsilon([A])$ in $\pi_1(\mathcal{P})$.*

Proof. The proof is essentially the same as in the case of exact categories (cf. [5, §4.9] or [7, Prop. 1.5(c)]).

Recall that by definition $\varepsilon([A]) = \psi_{[A],[A]}$ in $\pi_1(\mathcal{P})$. If $d : A \oplus A \rightarrow A \oplus A$ is the isomorphism $d(a, a') = (a', a)$ then it follows from the naturality and commutativity axioms that $\psi_{[A],[A]} = [d]$ in $\pi_1(\mathcal{P})$, and from the isomorphism of distinguished triangles

$$\begin{array}{ccccccc}
 A & \xrightarrow{(\text{id}, \text{id})} & A \oplus A & \xrightarrow{\langle \text{id}, -\text{id} \rangle} & A & \xrightarrow{0} & TA \\
 \downarrow \text{id} & & \downarrow d & & \downarrow -\text{id}_A & & \downarrow \text{id} \\
 A & \xrightarrow{(\text{id}, \text{id})} & A \oplus A & \xrightarrow{\langle \text{id}, -\text{id} \rangle} & A & \xrightarrow{0} & TA
 \end{array}$$

we deduce that $[d] = [-\text{id}_A]$ in $\pi_1(\mathcal{P})$. $\square$

Lemma 3.6. *If A is an object in $\mathcal{T}$ then $[A] \otimes [TA]$ has canonically the structure of a unit in $\mathcal{P}$, i.e. there exists a canonical isomorphism $\mu_A : [A] \otimes [TA] \rightarrow ([A] \otimes [TA]) \otimes ([A] \otimes [TA])$. This unit structure has the following properties.*

- (i) *If $a : A \rightarrow B$ is an isomorphism then $[a] \otimes [Ta]$ is an isomorphism of units $([A] \otimes [TA], \mu_A) \rightarrow ([B] \otimes [TB], \mu_B)$.*
- (ii) *If $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ is a distinguished triangle and Δ' denotes the distinguished triangle $TA \xrightarrow{Ta} TB \xrightarrow{Tb} TC \xrightarrow{-Tc} T^2A$, then the map*

$$\begin{aligned}
 [B] \otimes [TB] &\xrightarrow{[\Delta] \otimes [\Delta']} ([A] \otimes [C]) \otimes ([TA] \otimes [TC]) \\
 &\xrightarrow{\cong} ([A] \otimes [TA]) \otimes ([C] \otimes [TC])
 \end{aligned}$$

is an isomorphism of units $([B] \otimes [TB], \mu_B) \rightarrow ([A] \otimes [TA], \mu_A) \otimes ([C] \otimes [TC], \mu_C)$.

To simplify the notation we will always write μ_A for the natural isomorphisms $X \rightarrow ([A] \otimes [TA]) \otimes X$ and $X \rightarrow X \otimes ([A] \otimes [TA])$ associated to the unit $([A] \otimes [TA], \mu_A)$. The fact that $[A] \otimes [TA]$ has canonically the structure of a unit can also be expressed by saying that $[TA]$ is canonically a right inverse of $[A]$.

Proof of Lemma 3.6. From the distinguished triangle $\Delta_A : A \rightarrow 0 \rightarrow TA \xrightarrow{\text{id}} TA$ we obtain an isomorphism $[\Delta_A] : [0] \cong [A] \otimes [TA]$, and by Lemma 3.3 the object $[0]$ has a unit structure δ_0 . Hence $[A] \otimes [TA]$ has an induced unit structure $\mu_A = ([\Delta_A] \otimes [\Delta_A]) \circ \delta_0 \circ [\Delta_A]^{-1}$. Clearly μ_A is independent of the choice of zero object 0 .

Property (i) holds because an isomorphism $a : A \rightarrow B$ induces an isomorphism from the distinguished triangle Δ_A to Δ_B .

For property (ii) it suffices to show the commutativity of the following diagram.

$$\begin{array}{ccc}
[B] \otimes [TB] & \xrightarrow{[\Delta] \otimes [\Delta']} & ([A] \otimes [C]) \otimes ([TA] \otimes [TC]) \\
\downarrow [\Delta_B]^{-1} & & \downarrow \cong \\
[0] & & \\
\downarrow \delta_0 & & \\
[0] \otimes [0] & \xrightarrow{[\Delta_A] \otimes [\Delta_C]} & ([A] \otimes [TA]) \otimes ([C] \otimes [TC])
\end{array}$$

This diagram is the outer square in diagram (6) below. In (6), $\tilde{\Delta}$ denotes the distinguished triangle $C \xrightarrow{c} TA \xrightarrow{Ta} TB \xrightarrow{Tb} TC$ and $\rho_X : X \rightarrow X \otimes [0]$ is the natural isomorphism associated to the unit $[0]$.

$$\begin{array}{ccccc}
(6) \quad [B] \otimes [TB] & \xrightarrow{[\Delta] \otimes \text{id}} & [A] \otimes [C] \otimes [TB] & \xrightarrow{\text{id} \otimes \text{id} \otimes [\Delta']} & ([A] \otimes [C]) \otimes ([TA] \otimes [TC]) \\
\downarrow [\Delta_B]^{-1} & & \downarrow \text{id} \otimes [\tilde{\Delta}]^{-1} & & \downarrow \cong \\
[0] & \xrightarrow{[\Delta_A]} & [A] \otimes [TA] & \xrightarrow{\mu_C} & ([A] \otimes [TA]) \otimes ([C] \otimes [TC]) \\
\downarrow \rho_{[0]} = \delta_0 & & \downarrow \rho_{[A] \otimes [TA]} & & \parallel \\
[0] \otimes [0] & \xrightarrow{[\Delta_A] \otimes \text{id}} & ([A] \otimes [TA]) \otimes [0] & \xrightarrow{\text{id} \otimes \text{id} \otimes [\Delta_C]} & ([A] \otimes [TA]) \otimes ([C] \otimes [TC])
\end{array}$$

The commutativity of the top left hand square in (6) follows from the octahedral diagram

$$\begin{array}{ccccccc}
A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\
\parallel & & \downarrow & & \downarrow c & & \parallel \\
A & \longrightarrow & 0 & \longrightarrow & TA & \xrightarrow{\text{id}} & TA \\
& & \downarrow & & \downarrow Ta & & \\
& & TB & \xlongequal{\quad} & TB & & \\
& & \downarrow \text{id} & & \downarrow Tb & & \\
& & TB & \xrightarrow{Tb} & TC & &
\end{array}$$

The commutativity of the top right hand square in (6) follows from Lemma 3.8 below applied to $\tilde{\Delta}$ and its rotation Δ' (note that the proof of Lemma 3.8 does not use part (ii) of Lemma 3.6). The bottom left hand square in (6) is commutative by the naturality of ρ_X , and the bottom right hand square by definition of the unit structure on $[C] \otimes [TC]$. $\square$

Remark 3.7. Let 0 be a zero object in $\mathcal{T}$. Lemma 3.3 gives a unit structure on $[0]$ and on $[T0]$ and we can therefore consider the product unit structure on $[0] \otimes [T0]$. On the other hand Lemma 3.6 also gives a unit structure on $[0] \otimes [T0]$. It is not difficult to see that these two unit structures coincide.

Lemma 3.8. *Let $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ be a distinguished triangle and $\Delta' : B \xrightarrow{b} C \xrightarrow{c} TA \xrightarrow{-Ta} TB$ the rotated triangle. Then the composite isomorphism*

$$\begin{aligned} [B] &\xrightarrow{\mu_A} ([A] \otimes [TA]) \otimes [B] \\ &\xrightarrow{\cong} [A] \otimes ([B] \otimes [TA]) \\ &\xrightarrow{\text{id} \otimes [\Delta']^{-1}} [A] \otimes [C] \end{aligned}$$

is equal to $[\Delta] : [B] \rightarrow [A] \otimes [C]$.

Proof. Let $\alpha : [B] \rightarrow [A] \otimes [C]$ be the displayed isomorphism in the statement of the lemma. We must show that $[\Delta] \circ \alpha^{-1} = \text{id}_{[A] \otimes [C]}$. Consider the automorphism β of $[C]$ which is defined as the composite

$$[C] \xrightarrow{[\Delta']} [B] \otimes [TA] \xrightarrow{[\Delta] \otimes \text{id}} ([A] \otimes [C]) \otimes [TA] \cong ([A] \otimes [TA]) \otimes [C] \xrightarrow{\mu_A^{-1}} [C].$$

It is not hard to see that $\varepsilon([A]) \circ (\text{id}_{[A]} \otimes \beta) = [\Delta] \circ \alpha^{-1}$, therefore it suffices to show that $\beta = \varepsilon([A])$ in $\pi_1(\mathcal{P})$.

Consider the distinguished triangles

$$\begin{aligned} \Delta_1 : A &\xrightarrow{0} C \xrightarrow{(\text{id}, 0)} C \oplus TA \xrightarrow{\langle 0, \text{id} \rangle} TA, \\ \Delta_2 : C &\xrightarrow{(\text{id}, c)} C \oplus TA \xrightarrow{\langle c, -\text{id} \rangle} TA \xrightarrow{0} TC, \\ \Delta_3 : C &\xrightarrow{(\text{id}, 0)} C \oplus TA \xrightarrow{\langle 0, \text{id} \rangle} TA \xrightarrow{0} TC, \\ \Delta_4 : TA &\xrightarrow{(0, \text{id})} C \oplus TA \xrightarrow{\langle \text{id}, 0 \rangle} C \xrightarrow{0} T^2 A. \end{aligned}$$

The triangles Δ_2 and Δ_3 are isomorphic by the following diagram

$$(7) \quad \begin{array}{ccccccc} \Delta_3 : C & \xrightarrow{(\text{id}, 0)} & C \oplus TA & \xrightarrow{\langle 0, \text{id} \rangle} & TA & \xrightarrow{0} & TC \\ \downarrow \text{id} & & \downarrow d & & \downarrow \text{id} & & \downarrow \text{id} \\ \Delta_2 : C & \xrightarrow{(\text{id}, c)} & C \oplus TA & \xrightarrow{\langle c, -\text{id} \rangle} & TA & \xrightarrow{0} & TC, \end{array}$$

where $d : C \oplus TA \rightarrow C \oplus TA$ is the map $d(u, v) = (u, c(u) - v)$. We have $[d] = \varepsilon([A])$ in $\pi_1(\mathcal{P})$ because of the commutative diagram of distinguished triangles

$$\begin{array}{ccccccc} TA & \xrightarrow{(0, \text{id})} & C \oplus TA & \xrightarrow{\langle \text{id}, 0 \rangle} & C & \xrightarrow{0} & T^2 A \\ \downarrow -\text{id} & & \downarrow d & & \downarrow \text{id} & & \downarrow -\text{id} \\ TA & \xrightarrow{(0, \text{id})} & C \oplus TA & \xrightarrow{\langle \text{id}, 0 \rangle} & C & \xrightarrow{0} & T^2 A, \end{array}$$

Lemma 3.5, and the fact that $\varepsilon([A]) = \varepsilon([TA])$.

From the octahedral diagram

$$\begin{array}{ccccccc}
 A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\
 \parallel & & \downarrow b & & \downarrow (\text{id}, c) & & \parallel \\
 A & \xrightarrow{0} & C & \xrightarrow{(\text{id}, 0)} & C \oplus TA & \xrightarrow{\langle 0, \text{id} \rangle} & TA \\
 & & \downarrow c & & \downarrow \langle c, -\text{id} \rangle & & \\
 & & TA & \xlongequal{\quad} & TA & & \\
 & & \downarrow -Ta & & \downarrow 0 & & \\
 & & TB & \xrightarrow{Tb} & TC & &
 \end{array}$$

we obtain

$$([\Delta] \otimes \text{id}_{[TA]}) \circ [\Delta'] = \varphi \circ (\text{id}_{[A]} \otimes [\Delta_2]) \circ [\Delta_1].$$

Furthermore by (7) we have $[\Delta_2] = [\Delta_3] \circ \varepsilon([A])$, and from the commutativity axiom it follows that $[\Delta_4] = \psi \circ [\Delta_3]$. To show that $\beta = \varepsilon([A])$ it therefore suffices to show that

$$[C] \xrightarrow{[\Delta_1]} [A] \otimes [C \oplus TA] \xrightarrow{\text{id} \otimes [\Delta_4]} [A] \otimes ([TA] \otimes [C]) \cong ([A] \otimes [TA]) \otimes [C] \xrightarrow{\mu_A^{-1}} [C]$$

is the identity map. This follows easily from the octahedral diagram

$$\begin{array}{ccccccc}
 A & \longrightarrow & 0 & \longrightarrow & TA & \xrightarrow{\text{id}} & TA \\
 \parallel & & \downarrow & & \downarrow (0, \text{id}) & & \parallel \\
 A & \xrightarrow{0} & C & \xrightarrow{(\text{id}, 0)} & C \oplus TA & \xrightarrow{\langle 0, \text{id} \rangle} & TA \\
 & & \downarrow \text{id} & & \downarrow \langle \text{id}, 0 \rangle & & \\
 & & C & \xlongequal{\quad} & C & & \\
 & & \downarrow & & \downarrow 0 & & \\
 & & T0 & \longrightarrow & T^2 A & &
 \end{array}$$

and Lemma 3.4(ii) applied to the distinguished triangle $0 \rightarrow C \xrightarrow{\text{id}} C \rightarrow T0$. $\square$

Lemma 3.9. *Every commutative square*

$$\begin{array}{ccc}
 A' & \longrightarrow & B' \\
 \downarrow & & \downarrow \\
 A & \longrightarrow & B
 \end{array}$$

in $\mathcal{T}$ can be completed to a 9-term diagram

$$(8) \quad \begin{array}{ccccccc} A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & TA' \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & TA \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ A'' & \longrightarrow & B'' & \longrightarrow & C'' & \longrightarrow & TA'' \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ TA' & \dashrightarrow & TB' & \dashrightarrow & TC' & \dashrightarrow & T^2A' \end{array}$$

(i.e. the dashed arrows are the translates of the continuous arrows, all squares are commutative except for the bottom right hand square which is anti-commutative, the first three rows and the first three columns are distinguished triangles) such that for every determinant functor $[\cdot]$ on $\mathcal{T}$ with values in a Picard category $\mathcal{P}$ the induced diagram in $\mathcal{P}$

$$(9) \quad \begin{array}{ccc} [B] & \longrightarrow & [A] \otimes [C] \\ \downarrow & & \downarrow \\ & & ([A'] \otimes [A'']) \otimes ([C'] \otimes [C'']) \\ & & \downarrow \cong \\ [B'] \otimes [B''] & \longrightarrow & ([A'] \otimes [C']) \otimes ([A''] \otimes [C'']) \end{array}$$

is commutative.

Proof. In the proof of [1, Prop. 1.1.11] a 9-term diagram of the form (8) is constructed by combining three octahedral diagrams. The commutativity of diagram (9) then follows by applying the associativity axiom to these octahedral diagrams (using Lemma 3.8 to deal with the rotated triangles in the octahedral diagrams). We leave the details to the reader. $\square$

4. UNIVERSAL DETERMINANT FUNCTORS AND K -GROUPS

In this section we show that every small triangulated category $\mathcal{T}$ has a universal determinant functor. The groups $K_i(\mathcal{T})$ for $i = 0, 1$ are then defined as invariants of this universal determinant functor. The results in this section are motivated by Deligne's theory of universal determinant functors and virtual objects for exact categories which we recall first.

4.1. Universal determinant functors on exact categories. Let $\mathcal{E}$ be an exact category and w a class of morphisms in $\mathcal{E}$ as in §2.3.

Definition 4.1. A universal determinant functor on $(\mathcal{E}, w)$ is a Picard category $\mathcal{V}$ together with a determinant functor $f : (\mathcal{E}, w) \rightarrow \mathcal{V}$, such that for every Picard category $\mathcal{P}$ the functor

$$\mathrm{Hom}^\otimes(\mathcal{V}, \mathcal{P}) \rightarrow \det((\mathcal{E}, w), \mathcal{P})$$

which sends M to $M \circ f$ is an equivalence of categories.

If they exist, universal determinant functors are unique up to non-unique isomorphism, more precisely if $f : (\mathcal{E}, w) \rightarrow \mathcal{V}$ and $f' : (\mathcal{E}, w) \rightarrow \mathcal{V}'$ are universal determinant functors on $(\mathcal{E}, w)$ then there exists a monoidal functor $M : \mathcal{V} \rightarrow \mathcal{V}'$ (which is determined up to a non-unique isomorphism) such that the determinant functors $M \circ f$ and f' are (non-uniquely) isomorphic. Furthermore this functor $M : \mathcal{V} \rightarrow \mathcal{V}'$ is an equivalence of Picard categories.

Deligne showed in [5, §4] that there exists a universal determinant functor on every small exact category $\mathcal{E}$ (i.e. on $(\mathcal{E}, \text{iso})$). In fact he claims that the existence follows from “un argument standard”, however he does not indicate what this standard argument is. On the other hand Deligne also gives a topological construction of a universal determinant functor: using Quillen’s Q -construction he constructs a Picard category $V(\mathcal{E})$ (“catégorie des objets virtuels”) and a determinant functor $[\cdot] : \mathcal{E} \rightarrow V(\mathcal{E})$, and he then sketches a proof that it is universal. This construction shows that there exist canonical isomorphisms $\pi_i(V(\mathcal{E})) \cong K_i(\mathcal{E})$ for $i = 0, 1$, where $K_i(\mathcal{E})$ are the Quillen K -groups of $\mathcal{E}$.

In the case where $\mathcal{E}$ is the exact category of finitely generated projective modules over a ring, a different construction of a universal determinant functor on $\mathcal{E}$ was sketched by Fukaya and Kato in [6, §1.2]. However some details in their construction are omitted, and it is not obvious that one indeed obtains a determinant functor in the sense of Definition 2.4.

The proofs for the existence of a universal determinant functor by Deligne and by Fukaya and Kato are not very detailed. In §4.2 we will prove in detail the existence of a universal determinant functor on a triangulated category; it seems likely that a variant of that argument would also work for exact categories.

4.2. Universal determinant functors on triangulated categories. Let $\mathcal{T}$ be a triangulated category.

Definition 4.2. A universal determinant functor on $\mathcal{T}$ is a Picard category $\mathcal{V}$ together with a determinant functor $f : \mathcal{T} \rightarrow \mathcal{V}$, such that for every Picard category $\mathcal{P}$ the functor

$$\text{Hom}^{\otimes}(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$$

which sends M to $M \circ f$ is an equivalence of categories.

If they exist, universal determinant functors on triangulated categories are unique up to non-unique isomorphism, i.e. they satisfy the same uniqueness property as the universal determinant functors on exact categories in §4.1.

Theorem 4.3. *Let $\mathcal{T}$ be a small triangulated category. Then there exists a universal determinant functor $f : \mathcal{T} \rightarrow \mathcal{V}$.*

Proof. We proof the theorem by constructing $\mathcal{V}$ with generators and relations. Since $\mathcal{V}$ is a Picard category and therefore carries a lot of structure, the construction and the verification of the universal property are quite long and will occupy the remainder of §4.2.

Outline of the construction. Objects and morphisms of $\mathcal{V}$ are defined using words over certain alphabets. An alphabet $\mathbb{A}$ is a set and a word over $\mathbb{A}$ is a finite sequence of elements of $\mathbb{A}$. We denote the set of words over $\mathbb{A}$ by $\mathbb{A}^*$. If $A \in \mathbb{A}$ then we also write A for the word “ A ” consisting of the one element A . If $X, Y \in \mathbb{A}^*$ then XY denotes the concatenation of X and Y . For example in the construction

of the objects of $\mathcal{V}$ below, $\langle X \boxtimes Y \rangle$ denotes the word which is the concatenation of the words “ $\langle$ ”, X , “ $\boxtimes$ ”, Y and “ $\rangle$ ”.

The construction of the Picard category $\mathcal{V}$ proceeds in several steps. We first construct a set $\text{obj}(\mathcal{V})$ which will be the set of objects of $\mathcal{V}$. For every pair $X, Y \in \text{obj}(\mathcal{V})$ we then construct a set $\text{Hom}_{\mathcal{V}}(X, Y)$ which will be the set of morphisms from X to Y in $\mathcal{V}$. We define the composition of morphisms $\circ : \text{Hom}_{\mathcal{V}}(Y, Z) \times \text{Hom}_{\mathcal{V}}(X, Y) \rightarrow \text{Hom}_{\mathcal{V}}(X, Z)$ so that we obtain a category $\mathcal{V}$. Finally we define a tensor product $\otimes : \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$, an associativity constraint and a commutativity constraint which give an AC tensor structure on $\mathcal{V}$.

The most difficult step is the construction of the sets $\text{Hom}_{\mathcal{V}}(X, Y)$. The sets $\text{Hom}_{\mathcal{V}}(X, Y)$ are defined as $H(X, Y) / \sim_{X, Y}$ where $H(X, Y)$ is a set of words over an alphabet and $\sim_{X, Y}$ is an equivalence relation on $H(X, Y)$. To ensure that $\circ$ and $\otimes$ are defined on $\mathcal{V}$ and that we have a determinant functor $f : \mathcal{T} \rightarrow \mathcal{V}$ we must construct $H(X, Y)$ and $\sim_{X, Y}$ for all pairs X, Y simultaneously.

Constructing the objects. Let $\mathbb{A}$ be the alphabet

$$\mathbb{A} = \{\underline{A} : A \text{ is an object in } \mathcal{T}\} \cup \{\langle, \rangle, \boxtimes\}.$$

We define $\text{obj}(\mathcal{V})$ to be the smallest subset of $\mathbb{A}^*$ which satisfies the following two conditions.

- $\underline{A} \in \text{obj}(\mathcal{V})$ for every object A in $\mathcal{T}$.
- If $X, Y \in \text{obj}(\mathcal{V})$ then $\langle X \boxtimes Y \rangle \in \text{obj}(\mathcal{V})$.

If $X \in \text{obj}(\mathcal{V})$ then either $X = \underline{A}$ for some object A in $\mathcal{T}$, or $X = \langle Y \boxtimes Z \rangle$ for unique $Y, Z \in \text{obj}(\mathcal{V})$. This will allow us to argue by induction on the length of words.

Constructing the morphisms. Let $\mathbb{B}$ be the alphabet

$$\begin{aligned} \mathbb{B} = & \{\iota_X : X \in \text{obj}(\mathcal{V})\} \cup \{\langle, \rangle, \diamond, \boxtimes\} \\ & \cup \{\varphi_{X, Y, Z}, \overline{\varphi}_{X, Y, Z} : X, Y, Z \in \text{obj}(\mathcal{V})\} \cup \{\psi_{X, Y} : X, Y \in \text{obj}(\mathcal{V})\} \\ & \cup \{\underline{a} : a \text{ is an isomorphism in } \mathcal{T}\} \cup \{\underline{\Delta}, \overline{\Delta} : \Delta \text{ is a distinguished triangle in } \mathcal{T}\}. \end{aligned}$$

We define the system of sets $H(X, Y) \subseteq \mathbb{B}^*$ (parameterized by $X, Y \in \text{obj}(\mathcal{V})$) to be the smallest such system which satisfies the following nine conditions (where each condition must hold for all $X, Y, \dots \in \text{obj}(\mathcal{V})$).

- $\iota_X \in H(X, X)$.
- If $\alpha \in H(X, Y)$ and $\beta \in H(Y, Z)$ then $\langle \beta \diamond \alpha \rangle \in H(X, Z)$.
- If $\alpha_1 \in H(X_1, Y_1)$ and $\alpha_2 \in H(X_2, Y_2)$ then $\langle \alpha_1 \boxtimes \alpha_2 \rangle \in H(\langle X_1 \boxtimes X_2 \rangle, \langle Y_1 \boxtimes Y_2 \rangle)$.
- $\varphi_{X, Y, Z} \in H(\langle X \boxtimes \langle Y \boxtimes Z \rangle \rangle, \langle \langle X \boxtimes Y \rangle \boxtimes Z \rangle)$.
- $\overline{\varphi}_{X, Y, Z} \in H(\langle \langle X \boxtimes Y \rangle \boxtimes Z \rangle, \langle X \boxtimes \langle Y \boxtimes Z \rangle \rangle)$.
- $\psi_{X, Y} \in H(\langle X \boxtimes Y \rangle, \langle Y \boxtimes X \rangle)$.
- $\underline{a} \in H(\underline{A}, \underline{B})$ for every isomorphism $a : A \rightarrow B$ in $\mathcal{T}$.
- $\underline{\Delta} \in H(\underline{B}, \langle \underline{A} \boxtimes \underline{C} \rangle)$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$.
- $\overline{\Delta} \in H(\langle \underline{A} \boxtimes \underline{C} \rangle, \underline{B})$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$.

If $\alpha \in H(X, Y)$ then either α is one of $\iota_X, \varphi_{X, Y, Z}, \overline{\varphi}_{X, Y, Z}, \psi_{X, Y}, \underline{a}, \underline{\Delta}, \overline{\Delta}$, or $\alpha = \langle \gamma \diamond \beta \rangle$ for unique β and γ , or $\alpha = \langle \beta \boxtimes \gamma \rangle$ for unique β and γ . Hence we can argue

by induction on the length of words. Furthermore we observe that $H(U, V)$ and $H(U', V')$ are disjoint unless $U = U'$ and $V = V'$.

We want to define $\mathcal{V}$ to be the category whose objects are $\text{obj}(\mathcal{V})$ and whose morphisms from X to Y are $H(X, Y)/\sim_{X,Y}$, where $\sim_{X,Y}$ is an equivalence relation on $H(X, Y)$. The following definition formalizes the conditions we must impose on $\sim_{X,Y}$.

Definition 4.4. Suppose that for each pair $X, Y \in \text{obj}(\mathcal{V})$ we have an equivalence relation $\sim_{X,Y}$ on $H(X, Y)$. We say that such a system $\{\sim_{X,Y}\}$ is an admissible equivalence relation if it satisfies the following conditions (where each condition must hold for all $X, Y, \dots \in \text{obj}(\mathcal{V})$). To simplify the notation we omit the subscript X, Y of $\sim_{X,Y}$, and we do not explicitly mention in which set $H(X, Y)$ a relation $\alpha \sim \beta$ holds if it is clear from the context.

- (i) The following conditions ensure that $\mathcal{V}$ becomes a category with composition $\diamond$ and identity morphisms ι_X .
 - If $\alpha \sim \alpha'$ in $H(X, Y)$ and $\beta \sim \beta'$ in $H(Y, Z)$ then $\langle \beta \diamond \alpha \rangle \sim \langle \beta' \diamond \alpha' \rangle$.
 - $\langle \iota_Y \diamond \alpha \rangle \sim \alpha$ for all $\alpha \in H(X, Y)$.
 - $\langle \alpha \diamond \iota_X \rangle \sim \alpha$ for all $\alpha \in H(X, Y)$.
 - $\langle \gamma \diamond \langle \beta \diamond \alpha \rangle \rangle \sim \langle \langle \gamma \diamond \beta \rangle \diamond \alpha \rangle$ for all $\alpha \in H(X, Y)$, $\beta \in H(Y, Z)$, $\gamma \in H(Z, W)$.
- (ii) The following conditions ensure that $\boxtimes$ becomes a bifunctor on $\mathcal{V}$.
 - If $\alpha_1 \sim \alpha'_1$ in $H(X_1, Y_1)$ and $\alpha_2 \sim \alpha'_2$ in $H(X_2, Y_2)$ then $\langle \alpha_1 \boxtimes \alpha_2 \rangle \sim \langle \alpha'_1 \boxtimes \alpha'_2 \rangle$.
 - $\langle \iota_X \boxtimes \iota_Y \rangle \sim \iota_{\langle X \boxtimes Y \rangle}$.
 - If $\alpha_1 \in H(X_1, Y_1)$, $\beta_1 \in H(Y_1, Z_1)$, $\alpha_2 \in H(X_2, Y_2)$, $\beta_2 \in H(Y_2, Z_2)$ then $\langle \langle \beta_1 \diamond \alpha_1 \rangle \boxtimes \langle \beta_2 \diamond \alpha_2 \rangle \rangle \sim \langle \langle \beta_1 \boxtimes \beta_2 \rangle \diamond \langle \alpha_1 \boxtimes \alpha_2 \rangle \rangle$.
- (iii) The following conditions ensure that $\varphi_{X,Y,Z}$ and $\psi_{X,Y}$ become compatible associativity and commutativity constraints on $\mathcal{V}$.
 - ($\varphi_{X,Y,Z}$ is natural) If $\alpha \in H(X, X')$, $\beta \in H(Y, Y')$, $\gamma \in H(Z, Z')$ then $\langle \langle \langle \alpha \boxtimes \beta \rangle \boxtimes \gamma \rangle \diamond \varphi_{X,Y,Z} \rangle \sim \langle \varphi_{X',Y',Z'} \diamond \langle \alpha \boxtimes \langle \beta \boxtimes \gamma \rangle \rangle \rangle$.
 - ($\varphi_{X,Y,Z}$ is an isomorphism) $\langle \varphi_{X,Y,Z} \diamond \overline{\varphi}_{X,Y,Z} \rangle \sim \iota_{\langle \langle X \boxtimes Y \rangle \boxtimes Z \rangle}$ and $\langle \overline{\varphi}_{X,Y,Z} \diamond \varphi_{X,Y,Z} \rangle \sim \iota_{\langle X \boxtimes \langle Y \boxtimes Z \rangle \rangle}$.
 - (pentagonal axiom)

$$\begin{aligned} & \langle \varphi_{\langle X \boxtimes Y \rangle, Z, W} \diamond \varphi_{X, Y, \langle Z \boxtimes W \rangle} \rangle \\ & \sim \langle \langle \varphi_{X, Y, Z} \boxtimes \iota_W \rangle \diamond \langle \varphi_{X, Y \boxtimes Z, W} \diamond \langle \iota_X \boxtimes \varphi_{Y, Z, W} \rangle \rangle \rangle. \end{aligned}$$
 - ($\psi_{X,Y}$ is natural) If $\alpha \in H(X, X')$ and $\beta \in H(Y, Y')$ then $\langle \langle \beta \boxtimes \alpha \rangle \diamond \psi_{X,Y} \rangle \sim \langle \psi_{X',Y'} \diamond \langle \alpha \boxtimes \beta \rangle \rangle$.
 - $\langle \psi_{Y,X} \diamond \psi_{X,Y} \rangle \sim \iota_{\langle X \boxtimes Y \rangle}$.
 - (hexagonal axiom)

$$\begin{aligned} & \langle \varphi_{Z, X, Y} \diamond \langle \psi_{\langle X \boxtimes Y \rangle, Z} \diamond \varphi_{X, Y, Z} \rangle \rangle \\ & \sim \langle \langle \psi_{X, Z} \boxtimes \iota_Y \rangle \diamond \langle \varphi_{X, Z, Y} \diamond \langle \iota_X \boxtimes \psi_{Y, Z} \rangle \rangle \rangle. \end{aligned}$$
- (iv) The following conditions ensure that we obtain a functor $f_1 : \mathcal{T}_{\text{iso}} \rightarrow \mathcal{V}$ and isomorphisms $f_2(\Delta)$ in $\mathcal{V}$.
 - $\text{id}_A \sim \iota_A$ for every object A of $\mathcal{T}$.
 - $\underline{b \circ a} \sim \langle \underline{b} \diamond \underline{a} \rangle$ for all isomorphisms $a : A \rightarrow B$, $b : B \rightarrow C$ in $\mathcal{T}$.

- ($\underline{\Delta}$ is an isomorphism) $\langle \underline{\Delta} \diamond \overline{\Delta} \rangle \sim \iota_{\langle \underline{A} \boxtimes \underline{C} \rangle}$ and $\langle \overline{\Delta} \diamond \underline{\Delta} \rangle \sim \iota_{\underline{B}}$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$.
- (v) The following conditions ensure that the axioms for a determinant functor hold.
 - (naturality) $\langle \underline{\Delta}' \diamond \underline{b} \rangle \sim \langle \langle \underline{a} \boxtimes \underline{c} \rangle \diamond \underline{\Delta} \rangle$ for every isomorphism of distinguished triangles as in (4).
 - (associativity) $\langle \varphi_{\underline{A}, \underline{C}', \underline{A}'} \diamond \langle \langle \iota_{\underline{A}} \boxtimes \underline{\Delta}_{v2} \rangle \diamond \underline{\Delta}_{h2} \rangle \rangle \sim \langle \langle \underline{\Delta}_{h1} \boxtimes \iota_{\underline{A}'} \rangle \diamond \underline{\Delta}_{v1} \rangle$ for every octahedral diagram as in (3) (where Δ_{h1} and Δ_{h2} (resp. Δ_{v1} and Δ_{v2}) are the first and second horizontal (resp. vertical) distinguished triangles in diagram (3)).
 - (commutativity) $\langle \psi_{\underline{A}, \underline{B}} \diamond \underline{\Delta}_1 \rangle \sim \underline{\Delta}_2$ for every pair of distinguished triangles as in (5).

The intersection of admissible equivalence relations is again an admissible equivalence relation. Therefore there exists a unique minimal admissible equivalence relation which from now on we will denote by $\{\sim_{X,Y}\}$. For every pair of objects X, Y in $\mathcal{V}$ we define $\text{Hom}_{\mathcal{V}}(X, Y) = \text{H}(X, Y) / \sim_{X,Y}$. We denote the equivalence class of $\alpha \in \text{H}(X, Y)$ in $\text{Hom}_{\mathcal{V}}(X, Y) = \text{H}(X, Y) / \sim_{X,Y}$ by $[\alpha]$.

Constructing the Picard category. We define $\mathcal{V}$ to be the following category: the set of objects of $\mathcal{V}$ is $\text{obj}(\mathcal{V})$, the set of morphisms from $X \in \text{obj}(\mathcal{V})$ to $Y \in \text{obj}(\mathcal{V})$ is $\text{Hom}_{\mathcal{V}}(X, Y)$, and the composition of morphisms $[\alpha] \in \text{Hom}_{\mathcal{V}}(X, Y)$ and $[\beta] \in \text{Hom}_{\mathcal{V}}(Y, Z)$ is $[\beta] \circ [\alpha] = [\langle \beta \diamond \alpha \rangle] \in \text{Hom}_{\mathcal{V}}(X, Z)$. The definition of the equivalence relation ensures that this composition is well-defined and that we obtain a category. The identity morphism id_X of an object X is given by $\text{id}_X = [\iota_X]$.

Next we define a bifunctor $\otimes : \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$ by $X \otimes Y = \langle X \boxtimes Y \rangle$ for objects X, Y and $[\alpha] \otimes [\beta] = [\langle \alpha \boxtimes \beta \rangle]$ for morphisms $[\alpha], [\beta]$. Then $\varphi_{X,Y,Z} = [\varphi_{X,Y,Z}] \in \text{Hom}_{\mathcal{V}}(X \otimes (Y \otimes Z), (X \otimes Y) \otimes Z)$ and $\psi_{X,Y} = [\psi_{X,Y}] \in \text{Hom}_{\mathcal{V}}(X \otimes Y, Y \otimes X)$ give compatible associativity and commutativity constraints, so that $\mathcal{V}$ becomes an AC tensor category.

Lemma 4.5. *The AC tensor category $\mathcal{V}$ is a Picard category.*

Proof. Clearly $\mathcal{V}$ is non-empty. Also, since each of $[\iota_X]$, $[\varphi_{X,Y,Z}]$, $[\overline{\varphi}_{X,Y,Z}]$, $[\psi_{X,Y}]$, $[\underline{a}]$, $[\underline{\Delta}]$, $[\overline{\Delta}]$ is an isomorphism, it follows by induction that every morphism in $\mathcal{V}$ is an isomorphism.

Next we show that for every object X in $\mathcal{V}$ there exists an object A in $\mathcal{T}$ and an isomorphism $[\alpha] : X \rightarrow \underline{A}$ in $\mathcal{V}$ such that $\alpha \in \text{H}(X, \underline{A})$ is a word which does not contain any $\underline{\psi}$. If $X = \underline{A}$ we can take $\alpha = \iota_{\underline{A}}$. Now suppose that $X = \langle Y \boxtimes Z \rangle$ and that there exist isomorphisms $[\beta] : Y \rightarrow \underline{B}$ and $[\gamma] : Z \rightarrow \underline{C}$ where B and C are objects in $\mathcal{T}$ and $\beta \in \text{H}(Y, \underline{B})$ and $\gamma \in \text{H}(Z, \underline{C})$ do not contain any $\underline{\psi}$. Let Δ be a distinguished triangle of the form $B \rightarrow A \rightarrow C \rightarrow TB$. Then we can take $\alpha = \langle \overline{\Delta} \diamond \langle \beta \boxtimes \gamma \rangle \rangle \in \text{H}(X, \underline{A})$.

Now fix a zero object 0 in $\mathcal{T}$. We want to show that there exists a functorial isomorphism $\lambda_X : \underline{0} \otimes X \rightarrow X$ in $\mathcal{V}$. For $X = \underline{A}$ with A an object in $\mathcal{T}$ we let $\lambda_X = [\overline{\Delta}]$ where $\Delta : 0 \rightarrow A \xrightarrow{\text{id}} A \rightarrow T0$. For $X, Y \in \text{obj}(\mathcal{V})$ we define $\lambda_{\langle X \boxtimes Y \rangle}$ as the composite morphism

$$\underline{0} \otimes \langle X \boxtimes Y \rangle = \underline{0} \otimes (X \otimes Y) \xrightarrow{\varphi_{\underline{0}, X, Y}} (\underline{0} \otimes X) \otimes Y \xrightarrow{\lambda_X \otimes \text{id}_Y} X \otimes Y = \langle X \boxtimes Y \rangle.$$

To show that λ_X is functorial we must verify that

$$(10) \quad [\alpha] \circ \lambda_X = \lambda_Y \circ (\text{id}_0 \otimes [\alpha])$$

in $\mathcal{V}$ for all $\alpha \in H(X, Y)$. For this we use induction on the length of α . It is easy to check (10) for $\alpha = \iota_X$ and $\alpha = \underline{a}$. If $\alpha = \underline{\Delta}$ for a distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ then (10) follows from the associativity axiom applied to the octahedral diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{\text{id}} & A & \longrightarrow & T0 \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & B & \xrightarrow{\text{id}} & B & \longrightarrow & T0 \\ & & \downarrow & & \downarrow & & \\ & & C & \xlongequal{\quad} & C & & \\ & & \downarrow & & \downarrow & & \\ & & TA & \xrightarrow{\text{id}} & TA & & \end{array}$$

The statement also follows for $\alpha = \overline{\Delta}$ because $[\overline{\Delta}] = [\underline{\Delta}]^{-1}$ in $\mathcal{V}$. If $\alpha = \underline{\varphi}_{X,Y,Z}$ then we can deduce (10) from the pentagonal axiom and naturality of $\underline{\varphi}_{X,Y,Z}$, and if $\alpha = \overline{\varphi}_{X,Y,Z}$ then we use that $[\overline{\varphi}_{X,Y,Z}] = [\underline{\varphi}_{X,Y,Z}]^{-1}$ in $\mathcal{V}$. Next if (10) holds for α and β then it holds for $\langle \beta \diamond \alpha \rangle$ (if defined) and for $\langle \alpha \boxtimes \beta \rangle$. Hence it follows by induction that (10) is true for all α that do not contain any $\underline{\psi}$. Now consider $\alpha = \underline{\psi}_{X,Y}$. We have shown above that there exist objects A, B in $\mathcal{T}$ and isomorphisms $X \cong \underline{A}$, $Y \cong \underline{B}$ in $\mathcal{V}$ such that these isomorphisms are represented by words in $H(X, \underline{A})$ and $H(Y, \underline{B})$ which do not contain any $\underline{\psi}$. Hence for these isomorphisms we know (10) which by naturality of $\underline{\psi}$ reduces to the case $\alpha = \underline{\psi}_{\underline{A}, \underline{B}}$. In this case (10) follows from the commutativity axiom and the case of $\underline{\Delta}$ already shown above. Finally, using induction again we can deduce the validity of (10) for all $\alpha \in H(X, Y)$.

Now let $W \in \text{obj}(\mathcal{V})$. We must show that the functor $X \mapsto W \otimes X$ is an autoequivalence of $\mathcal{V}$. Since $W \cong \underline{A}$ for some object A in $\mathcal{T}$, it suffices to show this for the functor $F : X \mapsto \underline{A} \otimes X$. Let G be the functor $Y \mapsto \underline{TA} \otimes Y$. Then $G \circ F$ is isomorphic to the identity functor by

$$\underline{TA} \otimes (\underline{A} \otimes X) \cong (\underline{TA} \otimes \underline{A}) \otimes X \cong (\underline{A} \otimes \underline{TA}) \otimes X \xrightarrow{[\overline{\Delta}] \otimes \text{id}} \underline{0} \otimes X \xrightarrow{\lambda_X} X$$

where $\Delta : A \rightarrow 0 \rightarrow TA \xrightarrow{\text{id}} TA$, and $F \circ G$ is isomorphic to the identity functor by

$$\underline{A} \otimes (\underline{TA} \otimes X) \cong (\underline{A} \otimes \underline{TA}) \otimes X \xrightarrow{[\overline{\Delta}] \otimes \text{id}} \underline{0} \otimes X \xrightarrow{\lambda_X} X.$$

This completes the proof of Lemma 4.5. $\square$

Constructing the determinant functor. For an object A in $\mathcal{T}$ we define $f_1(A) = \underline{A} \in \text{obj}(\mathcal{V})$, and for an isomorphism $a : A \rightarrow B$ in $\mathcal{T}$ we define $f_1(a) = [\underline{a}] \in \text{Hom}_{\mathcal{V}}(f_1(A), f_1(B))$. For a distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ we define $f_2(\Delta) = [\underline{\Delta}] \in \text{Hom}_{\mathcal{V}}(f_1(B), f_1(A) \otimes f_1(C))$.

Lemma 4.6. $f = (f_1, f_2)$ is a determinant functor on $\mathcal{T}$ with values in $\mathcal{V}$.

Proof. This is immediate from the definition of the equivalence relation. $\square$

Verification of the universal property. We must show that for every Picard category $\mathcal{P}$ the functor $\text{Hom}^\otimes(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$, $M \mapsto M \circ f$, is an equivalence of categories.

Lemma 4.7. *The functor $\text{Hom}^\otimes(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$ is surjective on objects.*

Proof. Let $g = (g_1, g_2) : \mathcal{T} \rightarrow \mathcal{P}$ be a determinant functor. We want to show that there exists a monoidal functor $M : \mathcal{V} \rightarrow \mathcal{P}$ such that $g = M \circ f$.

On objects of $\mathcal{V}$ we define M by $M(\underline{A}) = g_1(A)$ and $M(\langle X \boxtimes Y \rangle) = M(X) \otimes M(Y)$. Next we define a morphism $M(\alpha) : M(X) \rightarrow M(Y)$ for every $\alpha \in H(X, Y)$. Firstly $M(\iota_X) = \text{id}_{M(X)}$. If $a : A \rightarrow B$ is an isomorphism in $\mathcal{T}$ then $M(\underline{a}) = g_1(a)$. If $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ is a distinguished triangle in $\mathcal{T}$ then $M(\underline{\Delta}) = g_2(\Delta)$ and $M(\overline{\Delta}) = g_2(\Delta)^{-1}$. Also $M(\varphi_{X,Y,Z}) = \varphi_{M(X), M(Y), M(Z)}$, $M(\overline{\varphi}_{X,Y,Z}) = \varphi_{M(X), M(Y), M(Z)}^{-1}$ and $M(\psi_{X,Y}) = \psi_{M(X), M(Y)}$. Finally $M(\langle \beta \diamond \alpha \rangle) = M(\beta) \circ M(\alpha)$ and $M(\langle \alpha_1 \boxtimes \alpha_2 \rangle) = M(\alpha_1) \otimes M(\alpha_2)$.

For a morphism $[\alpha] \in \text{Hom}_{\mathcal{V}}(X, Y) = H(X, Y) / \sim_{X,Y}$ we now define $M([\alpha]) = M(\alpha) \in \text{Hom}_{\mathcal{P}}(M(X), M(Y))$. To show that this is well-defined we must verify that the equivalence relation defined by $\alpha \sim \alpha'$ if $M(\alpha) = M(\alpha')$ is admissible. This is straightforward to check using that $\mathcal{P}$ is a Picard category and $g = (g_1, g_2)$ is a determinant functor.

Finally we define a natural isomorphism $c_{X,Y} : M(X) \otimes M(Y) \rightarrow M(X \otimes Y)$ by $c_{X,Y} = \text{id}_{M(X) \otimes M(Y)}$. Then it is easy to check that $M = (M, c)$ is a monoidal functor $\mathcal{V} \rightarrow \mathcal{P}$ and that $g = M \circ f$. $\square$

Lemma 4.8. *The functor $\text{Hom}^\otimes(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$ is fully faithful.*

Proof. Let $M = (M, c), M' = (M', c') : \mathcal{V} \rightarrow \mathcal{P}$ be two monoidal functors and let $\lambda : M \circ f \rightarrow M' \circ f$ be an isomorphism in $\det(\mathcal{T}, \mathcal{P})$. We must show that there exists a unique isomorphism $\kappa : M \rightarrow M'$ in $\text{Hom}^\otimes(\mathcal{V}, \mathcal{P})$ such that $\lambda = \kappa \circ f$.

If there is an isomorphism $\kappa : M \rightarrow M'$ such that $\lambda = \kappa \circ f$ then necessarily $\kappa_{\underline{A}} = \lambda_A$ for every object A in $\mathcal{T}$ and $\kappa_{\langle X \boxtimes Y \rangle} = (c'_{X,Y}) \circ (\kappa_X \otimes \kappa_Y) \circ c_{X,Y}^{-1}$ for all $X, Y \in \text{obj}(\mathcal{V})$. This shows that κ is uniquely determined by λ .

To show the existence of κ we define a morphism $\kappa_X : M(X) \rightarrow M'(X)$ for every object X in $\mathcal{V}$ as follows: $\kappa_{\underline{A}} = \lambda_A$ for every object A of $\mathcal{T}$, $\kappa_{\langle X \boxtimes Y \rangle} = (c'_{X,Y}) \circ (\kappa_X \otimes \kappa_Y) \circ c_{X,Y}^{-1}$. Using induction on the length of words which represent morphisms in $\mathcal{V}$, one can easily check that κ is an isomorphism of monoidal functors $M \rightarrow M'$, and by definition of κ it is obvious that $\lambda = \kappa \circ f$. $\square$

It follows from Lemmas 4.7 and 4.8 that for every Picard category $\mathcal{P}$ the functor $\text{Hom}^\otimes(\mathcal{V}, \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$ is an equivalence of categories. This completes the proof of Theorem 4.3. $\square$

4.3. K_0 and K_1 of a triangulated category.

Definition 4.9. Let $\mathcal{T}$ be a small triangulated category. For $i = 0, 1$ we define the group $K_i(\mathcal{T})$ by $K_i(\mathcal{T}) = \pi_i(\mathcal{V})$ where $f : \mathcal{T} \rightarrow \mathcal{V}$ is a universal determinant functor on $\mathcal{T}$.

The following argument shows that the groups $K_i(\mathcal{T})$ are well-defined and functorial in $\mathcal{T}$. Let $F : \mathcal{S} \rightarrow \mathcal{T}$ be an exact functor of small triangulated categories

and $f_S : \mathcal{S} \rightarrow \mathcal{V}(\mathcal{S})$, $f_T : \mathcal{T} \rightarrow \mathcal{V}(\mathcal{T})$ universal determinant functors for $\mathcal{S}$ and $\mathcal{T}$ respectively. The equivalence of categories $\text{Hom}^\otimes(\mathcal{V}(\mathcal{S}), \mathcal{V}(\mathcal{T})) \rightarrow \det(\mathcal{S}, \mathcal{V}(\mathcal{T}))$ shows that there exists a monoidal functor $M : \mathcal{V}(\mathcal{S}) \rightarrow \mathcal{V}(\mathcal{T})$, unique up to isomorphism, such that the determinant functors $M \circ f_S$ and $f_T \circ F$ in $\det(\mathcal{S}, \mathcal{V}(\mathcal{T}))$ are isomorphic. The monoidal functor M induces homomorphisms $\pi_i(M) : \pi_i(\mathcal{V}(\mathcal{S})) \rightarrow \pi_i(\mathcal{V}(\mathcal{T}))$ for $i = 0, 1$, and in fact these homomorphisms only depend on F because isomorphic monoidal functors induce the same maps on π_i . Applying this to the identity functor $F = \text{id} : \mathcal{T} \rightarrow \mathcal{T}$ implies that up to canonical isomorphism $K_i(\mathcal{T})$ is independent of the choice of universal determinant functor. For an arbitrary exact functor $F : \mathcal{S} \rightarrow \mathcal{T}$ one obtains an induced homomorphism $K_i(F) = \pi_i(M) : K_i(\mathcal{S}) \rightarrow K_i(\mathcal{T})$.

Lemma 4.10. *Let $F, F' : \mathcal{S} \rightarrow \mathcal{T}$ be exact functors of small triangulated categories. If F and F' are isomorphic then $K_i(F) = K_i(F') : K_i(\mathcal{S}) \rightarrow K_i(\mathcal{T})$ for $i = 0, 1$.*

Proof. Let $f_S : \mathcal{S} \rightarrow \mathcal{V}(\mathcal{S})$ and $f_T : \mathcal{T} \rightarrow \mathcal{V}(\mathcal{T})$ be universal determinant functors. Since F and F' are isomorphic exact functors it follows that $f_T \circ F$ and $f_T \circ F'$ are isomorphic determinant functors. Hence there exists a monoidal functor $M : \mathcal{V}(\mathcal{S}) \rightarrow \mathcal{V}(\mathcal{T})$ such that $M \circ f_S \cong f_T \circ F$ and $M \circ f_S \cong f_T \circ F'$. By definition $K_i(F) = \pi_i(M)$ and $K_i(F') = \pi_i(M)$. $\square$

This lemma immediately implies

Corollary 4.11. *If $F : \mathcal{S} \rightarrow \mathcal{T}$ is an equivalence of small triangulated categories then $K_i(F) : K_i(\mathcal{S}) \rightarrow K_i(\mathcal{T})$ is an isomorphism.*

Remark 4.12. Let $\mathcal{T}$ be a triangulated category such that the isomorphism classes of objects in $\mathcal{T}$ form a set. Then there exists a small full subcategory $\mathcal{S}$ of $\mathcal{T}$ satisfying $\mathcal{T}\mathcal{S} = \mathcal{S}$ and for which the inclusion functor $\mathcal{S} \rightarrow \mathcal{T}$ is an equivalence of categories. The triangulated category $\mathcal{T}$ induces the structure of a triangulated category on $\mathcal{S}$ and we can define $K_i(\mathcal{T})$ for $i = 0, 1$ by $K_i(\mathcal{T}) = K_i(\mathcal{S})$. Corollary 4.11 shows that up to canonical isomorphism the definition of $K_i(\mathcal{T})$ does not depend on the choice of $\mathcal{S}$.

Next we show that the group $K_0(\mathcal{T})$ is the Grothendieck group of the triangulated category $\mathcal{T}$. Recall that the Grothendieck group is defined to be the quotient $F(\mathcal{T})/R(\mathcal{T})$ where $F(\mathcal{T})$ is the free abelian group on isomorphism classes of objects of $\mathcal{T}$, and $R(\mathcal{T})$ is the subgroup of $F(\mathcal{T})$ which is generated by all elements $(B) - (A) - (C)$ whenever there is a distinguished triangle $A \rightarrow B \rightarrow C \rightarrow TA$ (here (A) denotes the element in $F(\mathcal{T})$ corresponding to the isomorphism class of the object A).

Proposition 4.13. *The group $K_0(\mathcal{T})$ is canonically isomorphic to the Grothendieck group of $\mathcal{T}$.*

Proof. Let $f = (f_1, f_2) : \mathcal{T} \rightarrow \mathcal{V}$ be a universal determinant functor for $\mathcal{T}$. We define a homomorphism $F(\mathcal{T}) \rightarrow K_0(\mathcal{T})$ by sending (A) to the isomorphism class of $f_1(A)$ in $\pi_0(\mathcal{V}) = K_0(\mathcal{T})$. One easily verifies that this homomorphism factors through a homomorphism $u : F(\mathcal{T})/R(\mathcal{T}) \rightarrow K_0(\mathcal{T})$.

Let $\mathcal{P}$ be the Picard category whose set of objects is the set $F(\mathcal{T})/R(\mathcal{T})$, whose morphisms are only the identity morphism of each object and whose tensor product is given by the product in the group $F(\mathcal{T})/R(\mathcal{T})$. There is a unique determinant functor $g : \mathcal{T} \rightarrow \mathcal{P}$ which sends an object A of $\mathcal{T}$ to the image of (A) in $F(\mathcal{T})/R(\mathcal{T})$. By the universality of $f : \mathcal{T} \rightarrow \mathcal{V}$ there exists a monoidal functor $M : \mathcal{V} \rightarrow \mathcal{P}$

(unique up to isomorphism) such that $M \circ f \cong g$, and by applying π_0 we obtain a homomorphism $v : K_0(\mathcal{T}) = \pi_0(\mathcal{V}) \xrightarrow{\pi_0(M)} \pi_0(\mathcal{P}) = F(\mathcal{T})/R(\mathcal{T})$.

It is not difficult to verify that $vu = \text{id}$ and that $uv(X) = X$ if $X \in \pi_0(\mathcal{V}) = K_0(\mathcal{T})$ is the isomorphism class of $f_1(A)$ for some object A in $\mathcal{T}$. But the proof of Theorem 4.3 shows that every object in $\mathcal{V}$ is isomorphic to $f_1(A)$ for some A , hence $uv = \text{id}$ follows. $\square$

We conclude this section with a description of the automorphisms of a determinant functor on $\mathcal{T}$ in terms of homomorphisms on $K_0(\mathcal{T})$.

Lemma 4.14. *Let $\mathcal{T}$ be a small triangulated category, $\mathcal{P}$ a Picard category and $g : \mathcal{T} \rightarrow \mathcal{P}$ a determinant functor. Then there exists a canonical isomorphism $\text{Aut}_{\det(\mathcal{T}, \mathcal{P})}(g) \cong \text{Hom}(K_0(\mathcal{T}), \pi_1(\mathcal{P}))$ (where Hom is homomorphisms of abelian groups).*

Proof. Let $f : \mathcal{T} \rightarrow \mathcal{V}$ be a universal determinant functor and $M : \mathcal{V} \rightarrow \mathcal{P}$ a monoidal functor for which $M \circ f \cong g$ in $\det(\mathcal{T}, \mathcal{P})$. Then $\text{Aut}_{\det(\mathcal{T}, \mathcal{P})}(g) \cong \text{Aut}_{\det(\mathcal{T}, \mathcal{P})}(M \circ f) \cong \text{Aut}_{\text{Hom}^\otimes(\mathcal{V}, \mathcal{P})}(M) \cong \text{Hom}(\pi_0(\mathcal{V}), \pi_1(\mathcal{P})) = \text{Hom}(K_0(\mathcal{T}), \pi_1(\mathcal{P}))$, where the second isomorphism follows from the universality of f and the third isomorphism is Lemma 2.3. $\square$

5. DETERMINANT FUNCTORS ON TRIANGULATED CATEGORIES WITH T-STRUCTURE

In this section we study the relation between the determinant functors on a triangulated category with a t-structure and the determinant functors on its heart.

5.1. Statement of result. Let $\mathcal{T}$ be a triangulated category with t-structure and let $\mathcal{E}$ be its heart. For the precise definition and a detailed discussion of these notions we refer the reader to [1, §1.3]. Here we just recall that a t-structure on a triangulated category $\mathcal{T}$ is a pair of full subcategories $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ of $\mathcal{T}$ satisfying various properties, and that the heart $\mathcal{E}$ of the t-structure is the intersection $\mathcal{E} = \mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq 0}$. The heart is a full abelian subcategory of $\mathcal{T}$. If $\Delta : 0 \rightarrow A \xrightarrow{a} B \xrightarrow{b} C \rightarrow 0$ is a short exact sequence in $\mathcal{E}$ then there exists a unique morphism $c : C \rightarrow TA$ in $\mathcal{T}$ such that $A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ is a distinguished triangle in $\mathcal{T}$. We will denote this distinguished triangle by $\widehat{\Delta}$.

Let $\mathcal{P}$ be a Picard category and $f = (f_1, f_2)$ a determinant functor on $\mathcal{T}$ with values in $\mathcal{P}$. We define a functor $g_1 : \mathcal{E}_{\text{iso}} \rightarrow \mathcal{P}$ to be the restriction of f_1 to $\mathcal{E}_{\text{iso}}$, and for every short exact sequence $\Delta : 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in $\mathcal{E}$ we define an isomorphism $g_2(\Delta) : g_1(B) \rightarrow g_1(A) \otimes g_1(C)$ by $g_2(\Delta) = f_2(\widehat{\Delta})$. Let $f|_{\mathcal{E}}$ denote the pair (g_1, g_2) .

Lemma 5.1. *If f is a determinant functor on the triangulated category $\mathcal{T}$ with values in $\mathcal{P}$, then $f|_{\mathcal{E}}$ is a determinant functor on the exact category $\mathcal{E}$ with values in $\mathcal{P}$. Moreover $f \mapsto f|_{\mathcal{E}}$ naturally extends to a functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$.*

Proof. If $\Delta \rightarrow \Delta'$ is an isomorphism of short exact sequences in $\mathcal{E}$ then we obtain an induced isomorphism of distinguished triangles $\widehat{\Delta} \rightarrow \widehat{\Delta}'$ in $\mathcal{T}$. Hence the naturality axiom for f implies the naturality axiom for $f|_{\mathcal{E}}$. Given a diagram of short exact sequences as in (1) then the associated diagram of distinguished triangles is an octahedral diagram, and therefore the associativity axiom for $f|_{\mathcal{E}}$ follows from the

associativity axiom for f . If Δ is the short exact sequence $A \rightarrow A \oplus B \rightarrow B$ then $\widehat{\Delta}$ is the distinguished triangle $A \rightarrow A \oplus B \rightarrow B \xrightarrow{0} TA$. Therefore the commutativity axiom for $f|_{\mathcal{E}}$ follows from the commutativity axiom for f .

We obtain a functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ because a morphism $\lambda : f \rightarrow f'$ in $\det(\mathcal{T}, \mathcal{P})$ restricts to a morphism $\lambda|_{\mathcal{E}} : f|_{\mathcal{E}} \rightarrow f'|_{\mathcal{E}}$ in $\det(\mathcal{E}, \mathcal{P})$. $\square$

Associated to the t-structure there is a canonical cohomological functor $H : \mathcal{T} \rightarrow \mathcal{E}$. The t-structure is called bounded if for every object A of $\mathcal{T}$ the cohomology objects $H(T^i A)$ are zero for all but finitely many $i \in \mathbb{Z}$. For any $n \in \mathbb{Z}$ we write $\mathcal{T}^{\leq n} = T^{-n}\mathcal{T}^{\leq 0}$ and $\mathcal{T}^{\geq n} = T^{-n}\mathcal{T}^{\geq 0}$. The t-structure is called non-degenerate if the intersections $\bigcap_{n \in \mathbb{Z}} \mathcal{T}^{\leq n}$ and $\bigcap_{n \in \mathbb{Z}} \mathcal{T}^{\geq n}$ consist only of zero objects. We can now state the main result of §5.

Theorem 5.2. *Let $\mathcal{T}$ be a triangulated category with a non-degenerate and bounded t-structure, and let $\mathcal{E}$ be its heart. Then for every Picard category $\mathcal{P}$ the restriction functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$, $f \mapsto f|_{\mathcal{E}}$, is an equivalence of categories.*

Corollary 5.3. *Let $\mathcal{T}$ be a small triangulated category with a non-degenerate and bounded t-structure, and let $\mathcal{E}$ be its heart. Then there exist canonical isomorphisms $K_i(\mathcal{T}) \cong K_i(\mathcal{E})$ for $i = 0, 1$.*

Proof. Fix universal determinant functors $f_{\mathcal{T}} : \mathcal{T} \rightarrow \mathcal{V}(\mathcal{T})$ and $f_{\mathcal{E}} : \mathcal{E} \rightarrow \mathcal{V}(\mathcal{E})$. Then by definition of a universal determinant functor there exists a monoidal functor $M : \mathcal{V}(\mathcal{E}) \rightarrow \mathcal{V}(\mathcal{T})$ (unique up to isomorphism) such that the determinant functors $(f_{\mathcal{T}})|_{\mathcal{E}}$ and $M \circ f_{\mathcal{E}}$ are isomorphic. For every Picard category $\mathcal{P}$ the functor $\text{Hom}^{\otimes}(\mathcal{V}(\mathcal{T}), \mathcal{P}) \rightarrow \text{Hom}^{\otimes}(\mathcal{V}(\mathcal{E}), \mathcal{P})$, $N \mapsto N \circ M$, is an equivalence of categories, because the diagram

$$\begin{array}{ccc} \text{Hom}^{\otimes}(\mathcal{V}(\mathcal{T}), \mathcal{P}) & \longrightarrow & \text{Hom}^{\otimes}(\mathcal{V}(\mathcal{E}), \mathcal{P}) \\ \downarrow & & \downarrow \\ \det(\mathcal{T}, \mathcal{P}) & \longrightarrow & \det(\mathcal{E}, \mathcal{P}) \end{array}$$

commutes up to natural isomorphism and the two vertical and the bottom horizontal functors are equivalences (by definition of a universal determinant functor and by Theorem 5.2 respectively). It follows that M is an equivalence of Picard categories and hence induces isomorphisms $K_i(\mathcal{E}) = \pi_i(\mathcal{V}(\mathcal{E})) \xrightarrow{\pi_i(M)} \pi_i(\mathcal{V}(\mathcal{T})) = K_i(\mathcal{T})$ for $i = 0, 1$. $\square$

As an important special case we note that if $\mathcal{E}$ is an abelian category and $\mathcal{T} = \mathcal{D}^b(\mathcal{E})$ its bounded derived category with the canonical t-structure then the assumptions of Corollary 5.3 are satisfied and hence $K_i(\mathcal{D}^b(\mathcal{E})) \cong K_i(\mathcal{E})$ for $i = 0, 1$.

To prove Theorem 5.2 we factorize the restriction functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ as the composite of two functors

$$(11) \quad \det(\mathcal{T}, \mathcal{P}) \xrightarrow{V} \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \xrightarrow{W} \det(\mathcal{E}, \mathcal{P})$$

and show that each of the functors V and W is an equivalence of categories.

The pair $(\text{Gr}^b(\mathcal{E}), T)$ consists of the category $\text{Gr}^b(\mathcal{E})$ of bounded $\mathbb{Z}$ -graded objects in $\mathcal{E}$ and the canonical translation functor T on $\text{Gr}^b(\mathcal{E})$. We will define determinant functors on exact categories with translation in §5.2, where we will also show that

for any exact category $\mathcal{E}$ the natural functor $W : \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ is an equivalence of categories. This deals with the functor W in (11).

Next in §5.3 we define the functor $V : \det(\mathcal{T}, \mathcal{P}) \rightarrow \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ for any triangulated category $\mathcal{T}$ with t-structure and heart $\mathcal{E}$, and we show that the restriction functor $\det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$, $f \mapsto f|_{\mathcal{E}}$, is the composite $W \circ V$.

In §5.4 we show that a bounded cohomological functor H from a triangulated category $\mathcal{T}$ to an abelian category $\mathcal{E}$ induces a functor

$$H^* : \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P}).$$

Finally in §5.5 we complete the proof of Theorem 5.2 by showing that if the t-structure on $\mathcal{T}$ is non-degenerate and bounded then V is an equivalence of categories. More precisely, we will show that under these assumptions the functor V is a quasi-inverse of the functor H^* where H is the canonical cohomological functor from $\mathcal{T}$ to its heart $\mathcal{E}$.

5.2. Exact categories with translation. If $\mathcal{E}$ is an exact category and T an exact automorphism of $\mathcal{E}$ then we will refer to T as a translation functor on $\mathcal{E}$ and to the pair $(\mathcal{E}, T)$ as an exact category with translation.

Definition 5.4. Let $(\mathcal{E}, T)$ be an exact category with translation and $\mathcal{P}$ a Picard category. A determinant functor on $(\mathcal{E}, T)$ with values in $\mathcal{P}$ is a triple (g_1, g_2, μ) , where (g_1, g_2) is a determinant functor on the exact category $\mathcal{E}$ with values in $\mathcal{P}$ and, for every object A in $\mathcal{E}$, μ_A is a unit structure on $g_1(A) \otimes g_1(TA)$, i.e. an isomorphism

$$\mu_A : g_1(A) \otimes g_1(TA) \rightarrow (g_1(A) \otimes g_1(TA)) \otimes (g_1(A) \otimes g_1(TA)),$$

such that the following two conditions hold.

- (i) If $a : A \rightarrow B$ is an isomorphism in $\mathcal{E}$ then $g_1(a) \otimes g_1(Ta)$ is an isomorphism of units $(g_1(A) \otimes g_1(TA), \mu_A) \rightarrow (g_1(B) \otimes g_1(TB), \mu_B)$.
- (ii) If $\Delta : 0 \rightarrow A \xrightarrow{a} B \xrightarrow{b} C \rightarrow 0$ is a short exact sequence in $\mathcal{E}$ and Δ' denotes the short exact sequence $0 \rightarrow TA \xrightarrow{Ta} TB \xrightarrow{Tb} TC \rightarrow 0$, then the composite map

$$\begin{aligned} g_1(B) \otimes g_1(TB) &\xrightarrow{g_2(\Delta) \otimes g_2(\Delta')} (g_1(A) \otimes g_1(C)) \otimes (g_1(TA) \otimes g_1(TC)) \\ &\xrightarrow{\cong} (g_1(A) \otimes g_1(TA)) \otimes (g_1(C) \otimes g_1(TC)) \end{aligned}$$

is an isomorphism of units $(g_1(B) \otimes g_1(TB), \mu_B) \rightarrow (g_1(A) \otimes g_1(TA), \mu_A) \otimes (g_1(C) \otimes g_1(TC), \mu_C)$.

A morphism $\lambda : (g_1, g_2, \mu) \rightarrow (g'_1, g'_2, \mu')$ of determinant functors on $(\mathcal{E}, T)$ with values in $\mathcal{P}$ is a morphism $\lambda : (g_1, g_2) \rightarrow (g'_1, g'_2)$ of the determinant functors on the underlying exact category which is compatible with the unit structures μ and μ' , i.e. for every object A in $\mathcal{E}$ the map $\lambda_A \otimes \lambda_{TA}$ is an isomorphism of units $(g_1(A) \otimes g_1(TA), \mu_A) \rightarrow (g'_1(A) \otimes g'_1(TA), \mu'_A)$.

We denote the category of determinant functors on $(\mathcal{E}, T)$ with values in $\mathcal{P}$ by $\det((\mathcal{E}, T), \mathcal{P})$.

If $\mathcal{E}$ is an exact category, then we write $\mathrm{Gr}^b(\mathcal{E})$ for the exact category whose objects are the bounded $\mathbb{Z}$ -graded objects in $\mathcal{E}$, i.e. tuples $(A_i)_{i \in \mathbb{Z}}$ of objects in $\mathcal{E}$ where A_i is a zero object for all but finitely many i , and morphisms $(A_i)_{i \in \mathbb{Z}} \rightarrow (B_i)_{i \in \mathbb{Z}}$ are tuples $(a_i)_{i \in \mathbb{Z}}$ where each $a_i : A_i \rightarrow B_i$ is a morphism in $\mathcal{E}$. The

category $\text{Gr}^b(\mathcal{E})$ has a canonical translation functor T given by $T((A_i)_{i \in \mathbb{Z}})_j = A_{j+1}$ for objects and $T((a_i)_{i \in \mathbb{Z}})_j = a_{j+1}$ for morphisms.

We embed $\mathcal{E}$ into $\text{Gr}^b(\mathcal{E})$ by sending objects (and morphisms) to tuples concentrated in degree 0. If (g_1, g_2, μ) is a determinant functor in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$ then the restriction of (g_1, g_2) to $\mathcal{E}$ is a determinant functor in $\det(\mathcal{E}, \mathcal{P})$. Clearly we can extend this to a functor $W : \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$.

Proposition 5.5. *The functor $W : \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ is an equivalence of categories.*

Proof. If $A = (A_i)$ is an object in $\text{Gr}^b(\mathcal{E})$ then we write A^{ev} and A^{od} for the objects $A^{\text{ev}} = \bigoplus_{i \text{ even}} A_i$ and $A^{\text{od}} = \bigoplus_{i \text{ odd}} A_i$ in $\mathcal{E}$. For a morphism $a = (a_i) : A \rightarrow B$ in $\text{Gr}^b(\mathcal{E})$ we obtain induced morphisms $a^{\text{ev}} : A^{\text{ev}} \rightarrow B^{\text{ev}}$ and $a^{\text{od}} : A^{\text{od}} \rightarrow B^{\text{od}}$. This defines two exact functors $\text{Gr}^b(\mathcal{E}) \rightarrow \mathcal{E}$.

For every object X in $\mathcal{P}$ we fix a right inverse X^* , i.e. an object X^* together with a unit structure on $X \otimes X^*$. If $\alpha : X \rightarrow Y$ is a morphism in $\mathcal{P}$ then we let $\alpha^* : X^* \rightarrow Y^*$ be the unique morphism for which $\alpha \otimes \alpha^* : X \otimes X^* \rightarrow Y \otimes Y^*$ is a morphism of units. Furthermore we note that there exists a canonical isomorphism $(X \otimes Y)^* \cong X^* \otimes Y^*$.

If (f_1, f_2) is a determinant functor in $\det(\mathcal{E}, \mathcal{P})$ then we define a functor $g_1 : \text{Gr}^b(\mathcal{E})_{\text{iso}} \rightarrow \mathcal{P}$ by $g_1(A) = f_1(A^{\text{ev}}) \otimes f_1(A^{\text{od}})^*$ for objects A in $\text{Gr}^b(\mathcal{E})$ and similarly for isomorphisms. For every short exact sequence $\Delta : A \rightarrow B \rightarrow C$ in $\text{Gr}^b(\mathcal{E})$ we define $g_2(\Delta) : g_1(B) \rightarrow g_1(A) \otimes g_1(C)$ to be the composite morphism

$$\begin{aligned} g_1(B) &= f_1(B^{\text{ev}}) \otimes f_1(B^{\text{od}})^* \\ &\xrightarrow{f_2(\Delta^{\text{ev}}) \otimes f_2(\Delta^{\text{od}})^*} (f_1(A^{\text{ev}}) \otimes f_1(C^{\text{ev}})) \otimes (f_1(A^{\text{od}}) \otimes f_1(C^{\text{od}}))^* \\ &\xrightarrow{\cong} (f_1(A^{\text{ev}}) \otimes f_1(C^{\text{ev}})) \otimes (f_1(A^{\text{od}})^* \otimes f_1(C^{\text{od}})^*) \\ &\xrightarrow{\cong} (f_1(A^{\text{ev}}) \otimes f_1(A^{\text{od}})^*) \otimes (f_1(C^{\text{ev}}) \otimes f_1(C^{\text{od}})^*) = g_1(A) \otimes g_1(C), \end{aligned}$$

where $\Delta^{\text{ev}} : A^{\text{ev}} \rightarrow B^{\text{ev}} \rightarrow C^{\text{ev}}$ and $\Delta^{\text{od}} : A^{\text{od}} \rightarrow B^{\text{od}} \rightarrow C^{\text{od}}$. Finally we define μ_A to be the unit structure on $g_1(A) \otimes g_1(TA)$ which is induced from the product unit structure on $(f_1(A^{\text{ev}}) \otimes f_1(A^{\text{ev}})^*) \otimes (f_1(A^{\text{od}}) \otimes f_1(A^{\text{od}})^*)$ via the isomorphism

$$\begin{aligned} g_1(A) \otimes g_1(TA) &= (f_1(A^{\text{ev}}) \otimes f_1(A^{\text{od}})^*) \otimes (f_1(A^{\text{od}}) \otimes f_1(A^{\text{ev}})^*) \\ &\xrightarrow{\cong} (f_1(A^{\text{ev}}) \otimes f_1(A^{\text{ev}})^*) \otimes (f_1(A^{\text{od}}) \otimes f_1(A^{\text{od}})^*) \\ &\xrightarrow{\varepsilon(f_1(A^{\text{od}}))} (f_1(A^{\text{ev}}) \otimes f_1(A^{\text{ev}})^*) \otimes (f_1(A^{\text{od}}) \otimes f_1(A^{\text{od}})^*). \end{aligned}$$

One can check that (g_1, g_2, μ) is a determinant functor in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$. Furthermore $(f_1, f_2) \mapsto (g_1, g_2, \mu)$ naturally extends to a functor $R : \det(\mathcal{E}, \mathcal{P}) \rightarrow \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$.

It is straightforward to verify that the composite functor

$$\det(\mathcal{E}, \mathcal{P}) \xrightarrow{R} \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \xrightarrow{W} \det(\mathcal{E}, \mathcal{P})$$

is isomorphic to the identity functor on $\det(\mathcal{E}, \mathcal{P})$.

Next we want to show that the composite of the functors

$$\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \xrightarrow{W} \det(\mathcal{E}, \mathcal{P}) \xrightarrow{R} \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$$

is isomorphic to the identity functor on $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$. For this we use the results from [2, §4]. Note that those results are formulated for determinant functors on certain categories of cochain complexes, but that they also apply to determinant functors on $(\mathrm{Gr}^b(\mathcal{E}), T)$. If (g_1, g_2, μ) is in $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ and $R(W(g_1, g_2, \mu)) = (g'_1, g'_2, \mu')$, then from [2, Prop. 4.4] we obtain an isomorphism of determinant functors $\pi : (g_1, g_2) \rightarrow (g'_1, g'_2)$ in $\det(\mathrm{Gr}^b(\mathcal{E}), \mathcal{P})$. Furthermore [2, Lemma 4.5] shows that π is compatible with μ and μ' and is therefore a morphism of determinant functors in $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$. Finally it is not difficult to verify that π is natural in g and hence gives a natural isomorphism from the identity functor on $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ to $R \circ W$. $\square$

5.3. Restriction functor to the graded category. Let $\mathcal{T}$ be a triangulated category with a t-structure and let $\mathcal{E}$ denote its heart. Let $f = (f_1, f_2) : \mathcal{T} \rightarrow \mathcal{P}$ be a determinant functor on $\mathcal{T}$. We want to show that f induces a determinant functor $V(f)$ on the exact category with translation $(\mathrm{Gr}^b(\mathcal{E}), T)$.

First we define a functor $J : \mathrm{Gr}^b(\mathcal{E}) \rightarrow \mathcal{T}$ as follows. If $A = (A_i)_{i \in \mathbb{Z}}$ is an object in $\mathrm{Gr}^b(\mathcal{E})$ then $J(A) = \bigoplus_{i \in \mathbb{Z}} T^{-i} A_i$ and if $a = (a_i)_{i \in \mathbb{Z}} : A \rightarrow B$ is a morphism in $\mathrm{Gr}^b(\mathcal{E})$ then $J(a) = \bigoplus_{i \in \mathbb{Z}} T^{-i} a_i : J(A) \rightarrow J(B)$. Note that there exists a canonical natural isomorphism $TJ \cong JT$. Let $\Delta : 0 \rightarrow A \xrightarrow{a} B \xrightarrow{b} C \rightarrow 0$ be a short exact sequence in $\mathrm{Gr}^b(\mathcal{E})$. Then for every $i \in \mathbb{Z}$ we have a short exact sequence $0 \rightarrow A_i \xrightarrow{a_i} B_i \xrightarrow{b_i} C_i \rightarrow 0$ in $\mathcal{E}$ and hence there is a unique morphism $c_i : C_i \rightarrow TA_i$ in $\mathcal{T}$ such that $A_i \xrightarrow{a_i} B_i \xrightarrow{b_i} C_i \xrightarrow{c_i} TA_i$ is a distinguished triangle. We define $J(\Delta)$ to be the distinguished triangle $J(A) \xrightarrow{J(a)} J(B) \xrightarrow{J(b)} J(C) \xrightarrow{c'} TJ(A)$ where $c' = \bigoplus_{i \in \mathbb{Z}} (-1)^i T^{-i} c_i$.

We define a functor $g_1 : \mathrm{Gr}^b(\mathcal{E})_{\mathrm{iso}} \rightarrow \mathcal{P}$ by $g_1 = f_1 \circ J$, and for every short exact sequence $\Delta : 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in $\mathrm{Gr}^b(\mathcal{E})$ we define an isomorphism $g_2(\Delta) : g_1(B) \rightarrow g_1(A) \otimes g_1(C)$ by $g_2(\Delta) = f_2(J(\Delta))$. For every object A in $\mathrm{Gr}^b(\mathcal{E})$ we define a unit structure μ'_A on $g_1(A) \otimes g_1(TA)$ by $\mu'_A = \mu_{J(A)}$ where $\mu_{J(A)}$ is the unit structure on $f_1(J(A)) \otimes f_1(TJ(A))$ defined in Lemma 3.6. Let $V(f)$ denote the triple (g_1, g_2, μ') .

Lemma 5.6. *If f is a determinant functor on $\mathcal{T}$ with values in $\mathcal{P}$, then $V(f)$ is a determinant functor on $(\mathrm{Gr}^b(\mathcal{E}), T)$ with values in $\mathcal{P}$. Moreover $f \mapsto V(f)$ naturally extends to a functor $V : \det(\mathcal{T}, \mathcal{P}) \rightarrow \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$.*

Proof. To see that (g_1, g_2) is a determinant functor on the exact category $\mathrm{Gr}^b(\mathcal{E})$ one proceeds as in the proof of Lemma 5.1. The necessary properties of the unit structure μ' follow from Lemma 3.6. The extension of $f \mapsto V(f)$ to a functor $V : \det(\mathcal{T}, \mathcal{P}) \rightarrow \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ is again similar to the proof of Lemma 5.1. $\square$

Lemma 5.7. *The composite functor $W \circ V : \det(\mathcal{T}, \mathcal{P}) \rightarrow \det(\mathcal{E}, \mathcal{P})$ is isomorphic to the restriction functor $f \mapsto f|_{\mathcal{E}}$.*

Proof. This is immediate from the definitions. $\square$

5.4. Cohomological functors. Let $\mathcal{T}$ be a triangulated category, $\mathcal{E}$ an abelian category and $H : \mathcal{T} \rightarrow \mathcal{E}$ a cohomological functor, i.e. an additive functor such that

for every distinguished triangle $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ in $\mathcal{T}$ the long sequence

$$(12) \quad \begin{aligned} \dots & \xrightarrow{H(T^{i-1}c)} H(T^i A) \xrightarrow{H(T^i a)} H(T^i B) \xrightarrow{H(T^i b)} H(T^i C) \\ & \xrightarrow{H(T^i c)} H(T^{i+1} A) \xrightarrow{H(T^{i+1} a)} \dots \end{aligned}$$

is exact in $\mathcal{E}$. We assume that H is bounded, i.e. that for every object A of $\mathcal{T}$ we have $H(T^i A) = 0$ for all but finitely many $i \in \mathbb{Z}$. Then we can define a functor $\tilde{H} : \mathcal{T} \rightarrow \text{Gr}^b(\mathcal{E})$ by setting $\tilde{H}(A) = (H(T^i A))_{i \in \mathbb{Z}}$ (resp. $\tilde{H}(a) = (H(T^i a))_{i \in \mathbb{Z}}$) for objects A (resp. morphisms a) in $\mathcal{T}$. The long exact sequence (12) can be written as the exact sequence

$$(13) \quad 0 \rightarrow \ker \tilde{H}a \rightarrow \tilde{H}A \xrightarrow{\tilde{H}a} \tilde{H}B \xrightarrow{\tilde{H}b} \tilde{H}C \xrightarrow{\tilde{H}c} T \ker \tilde{H}a \rightarrow 0$$

in $\text{Gr}^b(\mathcal{E})$.

Now consider a determinant functor $g = (g_1, g_2, \mu) : (\text{Gr}^b(\mathcal{E}), T) \rightarrow \mathcal{P}$ on the exact category with translation $(\text{Gr}^b(\mathcal{E}), T)$. We define a functor $f_1 : \mathcal{T}_{\text{iso}} \rightarrow \mathcal{P}$ as the composite $f_1 = g_1 \circ \tilde{H}$. Let $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ be a distinguished triangle in $\mathcal{T}$ and write the long exact cohomology sequence of Δ as the exact sequence (13) in $\text{Gr}^b(\mathcal{E})$. We define $f_2(\Delta) : f_1(B) \rightarrow f_1(A) \otimes f_1(C)$ to be the composite

$$\begin{aligned} f_1(B) = g_1(\tilde{H}B) & \xrightarrow{\mu_{\ker \tilde{H}a}} g_1(\tilde{H}B) \otimes g_1(\ker \tilde{H}a) \otimes g_1(T \ker \tilde{H}a) \\ & \xrightarrow{g_2(\Delta_1) \otimes \text{id} \otimes \text{id}} g_1(\ker \tilde{H}b) \otimes g_1(\ker \tilde{H}c) \otimes g_1(\ker \tilde{H}a) \otimes g_1(T \ker \tilde{H}a) \\ & \xrightarrow{\cong} g_1(\ker \tilde{H}a) \otimes g_1(\ker \tilde{H}b) \otimes g_1(\ker \tilde{H}c) \otimes g_1(T \ker \tilde{H}a) \\ & \xrightarrow{g_2(\Delta_2)^{-1} \otimes g_2(\Delta_3)^{-1}} g_1(\tilde{H}A) \otimes g_1(\tilde{H}C) = f_1(A) \otimes f_1(C), \end{aligned}$$

where the short exact sequence $\Delta_1 : 0 \rightarrow \ker \tilde{H}b \rightarrow \tilde{H}B \rightarrow \ker \tilde{H}c \rightarrow 0$, $\Delta_2 : 0 \rightarrow \ker \tilde{H}a \rightarrow \tilde{H}A \rightarrow \ker \tilde{H}b \rightarrow 0$ and $\Delta_3 : 0 \rightarrow \ker \tilde{H}c \rightarrow \tilde{H}C \rightarrow T \ker \tilde{H}a \rightarrow 0$ are obtained by splitting (13). Let $H^*(g)$ denote the pair (f_1, f_2) .

Proposition 5.8. *If g is a determinant functor in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$ then $H^*(g)$ is a determinant functor in $\det(\mathcal{T}, \mathcal{P})$. Moreover $g \mapsto H^*(g)$ naturally extends to a functor $H^* : \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$.*

Remark 5.9. The definition of $H^*(g)$ and Proposition 5.8 are also valid for an unbounded cohomological functor H if we use the category $\text{Gr}(\mathcal{E})$ of all $\mathbb{Z}$ -graded objects in $\mathcal{E}$ instead of $\text{Gr}^b(\mathcal{E})$.

Proof of Proposition 5.8. The second statement of the proposition follows easily once we have proved the first statement. Indeed, let $g = (g_1, g_2, \mu)$ and $g' = (g'_1, g'_2, \mu')$ be in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$ and define $H^*(g) = (f_1, f_2)$ and $H^*(g') = (f'_1, f'_2)$ as above. If $\lambda : g \rightarrow g'$ is a morphism in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$ then for every object A in $\mathcal{T}$ we can define $H^*(\lambda)_A = \lambda_{\tilde{H}A} : f_1(A) = g_1(\tilde{H}A) \rightarrow g'_1(\tilde{H}A) = f'_1(A)$. It is straightforward to check that $H^*(\lambda)$ is a natural isomorphism $f_1 \rightarrow f'_1$ and that $(H^*(\lambda)_A \otimes H^*(\lambda)_C) \circ f_2(\Delta) = f'_2(\Delta) \circ H^*(\lambda)_B$ for every distinguished triangle $\Delta : A \rightarrow B \rightarrow C \rightarrow TA$ in $\mathcal{T}$. Hence if we know that $H^*(g)$ and $H^*(g')$ are determinant functors, then $H^*(\lambda) : H^*(g) \rightarrow H^*(g')$ is a morphism of determinant functors, and we therefore obtain a functor $H^* : \det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P}) \rightarrow \det(\mathcal{T}, \mathcal{P})$.

The proof of the first statement of Proposition 5.8 will occupy the remainder of §5.4. We let $g = (g_1, g_2, \mu)$ be a determinant functor in $\det((\text{Gr}^b(\mathcal{E}), T), \mathcal{P})$ and

define $H^*(g) = (f_1, f_2)$ as above. To simplify the notation we will write H instead of $\tilde{H}$ for the rest of this proof.

An isomorphism of distinguished triangles induces an isomorphism of long exact cohomology sequences. From this it follows that f_2 is natural in isomorphisms of distinguished triangles.

The cohomology sequence (13) associated to a distinguished triangle Δ of the form $A \rightarrow A \oplus B \rightarrow B \xrightarrow{0} TA$ is of the form $0 \rightarrow 0 \rightarrow HA \rightarrow H(A \oplus B) \rightarrow HB \rightarrow T0 \rightarrow 0$. It follows easily that $f_2(\Delta) = g_2(\Delta')$ where Δ' is the short exact sequence $HA \rightarrow H(A \oplus B) \rightarrow HB$ in $\text{Gr}^b(\mathcal{E})$. Since $H(A \oplus B)$ is a direct sum of HA and HB and the maps $HA \rightarrow H(A \oplus B)$ and $H(A \oplus B) \rightarrow HB$ are the canonical inclusion and projection, we can deduce the commutativity axiom for $H^*(g)$ from the commutativity axiom for g .

It remains to show the associativity axiom. So assume that we are given the following octahedral diagram in $\mathcal{T}$.

$$(14) \quad \begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C' & \xrightarrow{fd} & TA \\ \parallel & & \downarrow c & & \downarrow d & & \parallel \\ A & \xrightarrow{ca} & C & \xrightarrow{e} & B' & \xrightarrow{f} & TA \\ & & \downarrow ge & & \downarrow g & & \\ & & A' & \xlongequal{\quad} & A' & & \\ & & \downarrow h & & \downarrow Tboh & & \\ & & TB & \xrightarrow{Tb} & TC' & & \end{array}$$

We must show that

$$\begin{array}{ccc} f_1(C) & \xrightarrow{f_2(\Delta_{h2})} & f_1(A) \otimes f_1(B') \\ \downarrow f_2(\Delta_{v1}) & & \downarrow \text{id} \otimes f_2(\Delta_{v2}) \\ f_1(B) \otimes f_1(A') & \xrightarrow{f_2(\Delta_{h1}) \otimes \text{id}} & f_1(A) \otimes f_1(C') \otimes f_1(A') \end{array}$$

is commutative in $\mathcal{P}$, where Δ_{h1} and Δ_{h2} (resp. Δ_{v1} and Δ_{v2}) are the first and second horizontal (resp. vertical) distinguished triangles in (14). By definition of f_1 and f_2 this is equivalent to showing that

$$(15) \quad \begin{array}{ccc} g_1(HC) & \xrightarrow{\mu_{\ker Hca}, g_2(\Delta_1), g_2(\Delta_2)^{-1}, g_2(\Delta_3)^{-1}} & g_1(HA) \otimes g_1(HB') \\ \downarrow \mu_{\ker Hc}, g_2(\Delta_4), g_2(\Delta_5)^{-1}, g_2(\Delta_6)^{-1} & & \downarrow \mu_{\ker Hd}, g_2(\Delta_7), g_2(\Delta_8)^{-1}, g_2(\Delta_9)^{-1} \\ g_1(HB) \otimes g_1(HA') & \xrightarrow{\mu_{\ker Ha}, g_2(\Delta_{10}), g_2(\Delta_{11})^{-1}, g_2(\Delta_{12})^{-1}} & g_1(HA) \otimes g_1(HC') \otimes g_1(HA') \end{array}$$

commutes in $\mathcal{P}$, where the short exact sequences $\Delta_1, \dots, \Delta_{12}$ are obtained by splitting the exact cohomology sequences associated to the distinguished triangles in

(14). To show the commutativity of (15) we must construct a huge commutative diagram in $\mathcal{P}$, which we do in several pieces.

First we observe that applying H to diagram (14) gives the following commutative diagram with exact rows and columns in $\text{Gr}^b(\mathcal{E})$

$$\begin{array}{ccccccccccc}
 & & & & 0 & & 0 & & & & \\
 & & & & \downarrow & & \downarrow & & & & \\
 & & & & \ker Hc & \longrightarrow & \ker Hd & & & & \\
 & & & & \downarrow & & \downarrow & & & & \\
 0 & \longrightarrow & \ker Ha & \longrightarrow & HA & \xrightarrow{Ha} & HB & \xrightarrow{Hb} & HC' & \xrightarrow{Hfd} & T\ker Ha \longrightarrow 0 \\
 & & \downarrow & & \parallel & & \downarrow Hc & & \downarrow Hd & & \downarrow \\
 0 & \longrightarrow & \ker Hca & \longrightarrow & HA & \xrightarrow{Hca} & HC & \xrightarrow{He} & HB' & \xrightarrow{Hf} & T\ker Hca \longrightarrow 0 \\
 & & & & & & \downarrow Hge & & \downarrow Hg & & \\
 & & & & & & HA' & \xlongequal{\quad} & HA' & & \\
 & & & & & & \downarrow Hh & & \downarrow H(Tb \circ h) & & \\
 & & & & & & T\ker Hc & \xrightarrow{HTb} & T\ker Hd & & \\
 & & & & & & \downarrow & & \downarrow & & \\
 & & & & & & 0 & & 0 & &
 \end{array}$$

and that applying H to the equality $Ta \circ f = h \circ g$ gives

$$(16) \quad THa \circ Hf = Hh \circ Hg.$$

In $\text{Gr}^b(\mathcal{E})$ we have the following short exact sequences.

$$\begin{aligned}
 \Delta_1 : \ker He &\rightarrow HC \xrightarrow{He} \ker Hf \\
 \Delta_2 : \ker Hca &\rightarrow HA \xrightarrow{Hca} \ker He \\
 \Delta_3 : \ker Hf &\rightarrow HB' \xrightarrow{Hf} T\ker Hca \\
 \Delta_4 : \ker Hge &\rightarrow HC \xrightarrow{Hge} \ker Hh \\
 \Delta_5 : \ker Hc &\rightarrow HB \xrightarrow{Hc} \ker Hge \\
 \Delta_6 : \ker Hh &\rightarrow HA' \xrightarrow{Hh} T\ker Hc \\
 \Delta_7 : \ker Hg &\rightarrow HB' \xrightarrow{Hg} \ker H(Tb \circ h) \\
 \Delta_8 : \ker Hd &\rightarrow HC' \xrightarrow{Hd} \ker Hg \\
 \Delta_9 : \ker H(Tb \circ h) &\rightarrow HA' \xrightarrow{H(Tb \circ h)} T\ker Hd
 \end{aligned}$$

$$\begin{aligned}
\Delta_{10} &: \ker Hb \rightarrow HB \xrightarrow{Hb} \ker Hfd \\
\Delta_{11} &: \ker Ha \rightarrow HA \xrightarrow{Ha} \ker Hb \\
\Delta_{12} &: \ker Hfd \rightarrow HC' \xrightarrow{Hfd} T\ker Ha \\
\Delta_{13} &: \ker Ha \rightarrow \ker Hca \xrightarrow{Ha} \ker Hb \cap \ker Hc \\
\Delta_{14} &: \ker Hb \cap \ker Hc \rightarrow \ker Hc \xrightarrow{Hb} \ker Hd
\end{aligned}$$

Furthermore we will need the translations of Δ_{13} and Δ_{14} , that is the following two short exact sequences.

$$\begin{aligned}
T\Delta_{13} &: T\ker Ha \rightarrow T\ker Hca \xrightarrow{THa} T\ker Hb \cap T\ker Hc \\
T\Delta_{14} &: T\ker Hb \cap T\ker Hc \rightarrow T\ker Hc \xrightarrow{THb} T\ker Hd
\end{aligned}$$

We will show the commutativity of the following diagram. Since its outer square is (15) this will imply the associativity axiom for $H^*(g)$. For typographic reasons we will write $[\cdot]$ for $g_1(\cdot)$ and omit the word $\ker$ in all diagrams below, so for example we have $[HA] = g_1(HA)$ and $[Ha] = g_1(\ker Ha)$. Furthermore we will write Δ_i for both $g_2(\Delta_i)$ and $g_2(\Delta_i)^{-1}$.

$$\begin{array}{ccccc}
[HC] & \xrightarrow{\mu_{Hca}} & [HC] & \xrightarrow{\Delta_1, \Delta_2, \Delta_3} & [HA] \otimes [HB'] \\
\downarrow \mu_{Hc} & \searrow \mu_{Hd}, \mu_{Ha}, \mu_{Hb \cap Hc} & \downarrow \mu_{Hd}, \Delta_{13}, T\Delta_{13} & & \downarrow \mu_{Hd} \\
[HC] & & [HC] & & [HA] \otimes [HB'] \\
\otimes [Hc] \otimes [THc] & \xrightarrow{\mu_{Ha}, \Delta_{14}, T\Delta_{14}} & \otimes [Ha] \otimes [THa] & \xrightarrow{\Delta_{13}, T\Delta_{13}, \Delta_1, \Delta_2, \Delta_3} & \otimes [Hd] \otimes [THd] \\
\downarrow \Delta_4, \Delta_5, \Delta_6 & & \downarrow \Delta_{14}, T\Delta_{14}, \Delta_4, \Delta_5, \Delta_6 & & \downarrow \Delta_7, \Delta_8, \Delta_9 \\
[HB] \otimes [HA'] & \xrightarrow{\mu_{Ha}} & [HB] \otimes [HA'] & \xrightarrow{\Delta_{10}, \Delta_{11}, \Delta_{12}} & [HA] \otimes [HC'] \otimes [HA'] \\
& & \otimes [Ha] \otimes [THa] & &
\end{array}$$

From the properties of μ it immediately follows that the two triangles are commutative. Also the commutativity of the top right and bottom left hand squares is obvious.

It remains to show that the bottom right hand square is commutative. This square can be subdivided as follows.

$$\begin{array}{ccccc}
\begin{array}{c} [HC] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow{\Delta_1} & \begin{array}{c} [He] \otimes [Hf] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow[\Delta_2, \Delta_3]{\Delta_{13}, T\Delta_{13},} & \begin{array}{c} [HA] \otimes [HB'] \\ \otimes [Hd] \otimes [THd] \end{array} \\
\downarrow \Delta_4 & S_1 & \downarrow \Delta_{15} & S_2 & \downarrow \Delta_7 \\
\begin{array}{c} [Hge] \otimes [Hh] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow{\Delta_{16}} & \begin{array}{c} [He] \otimes [Hf \cap Hg] \otimes [Hh] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow[\Delta_{17}, \Delta_{18}]{\Delta_2, \Delta_{13},} & \begin{array}{c} [HA] \\ \otimes [Hg] \otimes [H(Tb \circ h)] \\ \otimes [Hd] \otimes [THd] \end{array} \\
\downarrow \begin{array}{c} \Delta_{14}, T\Delta_{14}, \\ \Delta_5, \Delta_6 \end{array} & S_3 & \downarrow \begin{array}{c} \Delta_6, T\Delta_{14}, \\ \Delta_{19}, \Delta_{20} \end{array} & S_4 & \downarrow \Delta_8, \Delta_9 \\
\begin{array}{c} [HB] \otimes [HA'] \\ \otimes [Ha] \otimes [THa] \end{array} & \xrightarrow{\Delta_{10}} & \begin{array}{c} [Hb] \otimes [Hfd] \\ \otimes [Ha] \otimes [THa] \\ \otimes [HA'] \end{array} & \xrightarrow{\Delta_{11}, \Delta_{12}} & [HA] \otimes [HC'] \otimes [HA']
\end{array}$$

Here $\Delta_{15}, \dots, \Delta_{20}$ are the following short exact sequences in $\text{Gr}^b(\mathcal{E})$.

$$\begin{aligned}
\Delta_{15} &: \ker Hf \cap \ker Hg \rightarrow \ker Hf \xrightarrow{Hg} \ker Hh \\
\Delta_{16} &: \ker He \rightarrow \ker Hge \xrightarrow{He} \ker Hf \cap \ker Hg \\
\Delta_{17} &: \ker Hf \cap \ker Hg \rightarrow \ker Hg \xrightarrow{Hf} T \ker Ha \\
\Delta_{18} &: \ker Hh \rightarrow \ker H(Tb \circ h) \xrightarrow{Hh} T \ker Hb \cap T \ker Hc \\
\Delta_{19} &: \ker Hb \cap \ker Hc \rightarrow \ker Hb \xrightarrow{Hc} \ker He \\
\Delta_{20} &: \ker Hd \rightarrow \ker Hfd \xrightarrow{Hd} \ker Hf \cap \ker Hg
\end{aligned}$$

Applying the associativity axiom to the commutative diagram of short exact sequences

$$\begin{array}{ccccc}
\ker He & \longrightarrow & \ker Hge & \xrightarrow{He} & \ker Hf \cap \ker Hg \\
\parallel & & \downarrow & & \downarrow \\
\ker He & \longrightarrow & HC & \xrightarrow{He} & \ker Hf \\
& & \downarrow Hge & & \downarrow Hg \\
& & \ker Hh & \xlongequal{\quad} & \ker Hh
\end{array}$$

gives the commutativity of the square S_1 .

Next we will show the commutativity of the square S_2 . For this we subdivide S_2 as in the following diagram

$$\begin{array}{ccccc}
\begin{array}{l} [He] \otimes [Hf] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow{\Delta_{13}, \Delta_2, T\Delta_{13}, \Delta_3} & \begin{array}{l} [HA] \otimes [HB'] \\ \otimes [Hd] \otimes [THd] \end{array} \\
\downarrow \Delta_{15} & \searrow \Delta_{13}, \Delta_2, \Delta_{21} & \swarrow \Delta_{22} & \downarrow \Delta_7 \\
\begin{array}{l} [He] \otimes [Hf \cap Hg] \otimes [Hh] \\ \otimes [Ha] \otimes [THa] \\ \otimes [Hb \cap Hc] \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow{\Delta_{13}, \Delta_2, \Delta_{17}} & \begin{array}{l} [HA] \otimes [H(Ta \circ f)] \\ \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & \xrightarrow{\Delta_{18}} & \begin{array}{l} [HA] \otimes [H(Tb \circ h)] \\ \otimes [Hd] \otimes [THd] \end{array} \\
& & \downarrow \Delta_{23} & & \\
& & \begin{array}{l} [HA] \otimes [Hg] \otimes [Hh] \\ \otimes [THb \cap THc] \\ \otimes [Hd] \otimes [THd] \end{array} & &
\end{array}$$

$S_{2,1}$ (top triangle), $S_{2,2}$ (right triangle), $S_{2,3}$ (left triangle)

where $\Delta_{21}, \dots, \Delta_{23}$ are the short exact sequences

$$\begin{aligned}
\Delta_{21} : \ker Hf &\rightarrow \ker H(Ta \circ f) \xrightarrow{Hf} T \ker Ha \\
\Delta_{22} : \ker H(Ta \circ f) &\rightarrow HB' \xrightarrow{H(Ta \circ f)} T \ker Hb \cap T \ker Hc \\
\Delta_{23} : \ker Hg &\rightarrow \ker H(Ta \circ f) \xrightarrow{Hg} \ker Hh.
\end{aligned}$$

The commutativity of $S_{2,1}$ and $S_{2,2}$ comes from the associativity axiom applied to the following two commutative diagrams of short exact sequences

$$\begin{array}{ccccc}
\ker Hf & \longrightarrow & \ker H(Ta \circ f) & \xrightarrow{Hf} & T \ker Ha \\
\parallel & & \downarrow & & \downarrow \\
\ker Hf & \longrightarrow & HB' & \xrightarrow{Hf} & T \ker Hca \\
& & \downarrow H(Ta \circ f) & & \downarrow THa \\
& & T \ker Hb \cap T \ker Hc & \xlongequal{\quad} & T \ker Hb \cap T \ker Hc,
\end{array}$$

$$\begin{array}{ccccc}
\ker Hg & \longrightarrow & \ker H(Ta \circ f) & \xrightarrow{Hg} & \ker Hh \\
\parallel & & \downarrow & & \downarrow \\
\ker Hg & \longrightarrow & HB' & \xrightarrow{Hg} & \ker H(Tb \circ h) \\
& & \downarrow H(Ta \circ f) & & \downarrow Hh \\
& & T \ker Hb \cap T \ker Hc & \xlongequal{\quad} & T \ker Hb \cap T \ker Hc,
\end{array}$$

(where the last square in the second diagram is commutative by (16)), and the commutativity of $S_{2,3}$ comes from [7, Cor. 1.10] applied to

$$\begin{array}{ccccc}
 \ker Hf \cap \ker Hg & \longrightarrow & \ker Hg & \xrightarrow{Hf} & T \ker Ha \\
 \downarrow & & \downarrow & & \parallel \\
 \ker Hf & \longrightarrow & \ker H(Ta \circ f) & \xrightarrow{Hf} & T \ker Ha \\
 \downarrow Hg & & \downarrow Hg & & \\
 \ker Hh & \xlongequal{\quad} & \ker Hh & &
 \end{array}$$

The commutativity of the squares S_3 and S_4 can be shown similarly and is left to the reader. $\square$

5.5. Completion of the proof. Let $\mathcal{T}$ be a triangulated category with a bounded t-structure, let $\mathcal{E}$ be its heart and $H : \mathcal{T} \rightarrow \mathcal{E}$ the canonical cohomological functor. Recall that in §5.3 and §5.4 we defined functors

$$\begin{aligned}
 V : \det(\mathcal{T}, \mathcal{P}) &\rightarrow \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}), \\
 H^* : \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}) &\rightarrow \det(\mathcal{T}, \mathcal{P}).
 \end{aligned}$$

In this subsection we show that if the t-structure is non-degenerate and bounded then V is an equivalence of categories with quasi-inverse H^* .

Lemma 5.10. *Assume that the t-structure on $\mathcal{T}$ is bounded. Then the composite functor*

$$\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}) \xrightarrow{H^*} \det(\mathcal{T}, \mathcal{P}) \xrightarrow{V} \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$$

is isomorphic to the identity functor on $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$.

Proof. The composite functor $\mathrm{Gr}^b(\mathcal{E}) \xrightarrow{J} \mathcal{T} \xrightarrow{\bar{H}} \mathrm{Gr}^b(\mathcal{E})$ is canonically isomorphic to the identity functor on $\mathrm{Gr}^b(\mathcal{E})$. It follows that if $g = (g_1, g_2, \mu) \in \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ and $V(H^*(g)) = (g'_1, g'_2, \mu')$ then there exists a canonical natural isomorphism $\xi : g_1 \rightarrow g'_1$. It is easily verified that ξ is compatible with the isomorphisms $g_2(\Delta)$ and $g'_2(\Delta)$ for every short exact sequence Δ in $\mathrm{Gr}^b(\mathcal{E})$ and with the unit structures μ_A and μ'_A for every object A , and thus is an isomorphism of determinant functors $\xi : g \rightarrow V(H^*(g))$.

It is then straightforward to check that ξ is natural in g and hence gives a natural isomorphism from the identity functor on $\det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P})$ to $V \circ H^*$. $\square$

Proposition 5.11. *Assume that the t-structure on $\mathcal{T}$ is non-degenerate and bounded. Then the composite functor*

$$\det(\mathcal{T}, \mathcal{P}) \xrightarrow{V} \det((\mathrm{Gr}^b(\mathcal{E}), T), \mathcal{P}) \xrightarrow{H^*} \det(\mathcal{T}, \mathcal{P})$$

is isomorphic to the identity functor on $\det(\mathcal{T}, \mathcal{P})$.

Proof. We will show that for every determinant functor f in $\det(\mathcal{T}, \mathcal{P})$ there exists a canonical isomorphism of determinant functors $\eta : f \rightarrow H^*(V(f))$. To complete the proof of the proposition one must then check that η is natural in f . This follows almost immediately from the definition of η and is left to the reader.

Write $f = (f_1, f_2)$ and $H^*(V(f)) = (f'_1, f'_2)$. We want to construct a natural isomorphism $\eta : f_1 \rightarrow f'_1$ that is compatible with f_2 and f'_2 .

By definition of H^* and V we have $f'_1(A) = f_1(J\tilde{H}A)$ for an object A of $\mathcal{T}$ and similarly for isomorphisms. Since the t-structure is non-degenerate and bounded it follows from [1, Prop. 1.3.7] that $A \in \mathcal{T}^{\geq i}$ for some $i \in \mathbb{Z}$. Then $\tau_{\leq i}A \cong T^{-i}H^iA$ where we write $H^iA = HT^iA$. Applying f_2 to the distinguished triangle $\tau_{\leq i}A \rightarrow A \rightarrow \tau_{>i}A \rightarrow T\tau_{\leq i}A$ gives an isomorphism

$$\begin{aligned} f_1(A) &\cong f_1(\tau_{\leq i}A) \otimes f_1(\tau_{>i}A) \\ &\cong f_1(T^{-i}H^iA) \otimes f_1(\tau_{>i}A). \end{aligned}$$

Next we apply the same argument to $\tau_{>i}A \in \mathcal{T}^{\geq i+1}$ and obtain

$$\begin{aligned} f_1(\tau_{>i}A) &\cong f_1(T^{-(i+1)}H^{i+1}\tau_{>i}A) \otimes f_1(\tau_{>i+1}\tau_{>i}A) \\ &\cong f_1(T^{-(i+1)}H^{i+1}A) \otimes f_1(\tau_{>i+1}A), \end{aligned}$$

because $H^{i+1}\tau_{>i}A \cong H^{i+1}A$ and $\tau_{>i+1}\tau_{>i}A \cong \tau_{>i+1}A$. We then apply the same argument to $\tau_{>i+1}A \in \mathcal{T}^{\geq i+2}$, etc. By [1, Prop. 1.3.7] we have $\tau_{>i+j}A = 0$ for sufficiently large j , and therefore we obtain an isomorphism

$$\begin{aligned} \eta_A : f_1(A) &\cong \bigotimes_{i \in \mathbb{Z}} f_1(T^{-i}H^iA) \\ &\cong f_1\left(\bigoplus_{i \in \mathbb{Z}} T^{-i}H^iA\right) \\ &= f_1(J\tilde{H}A) = f'_1(A). \end{aligned}$$

Obviously the isomorphism η_A is functorial for isomorphisms $A \rightarrow B$.

To prove that η is an isomorphism of determinant functors, we must show that for every distinguished triangle $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ in $\mathcal{T}$ the diagram

$$(17) \quad \begin{array}{ccc} f_1(B) & \xrightarrow{f_2(\Delta)} & f_1(A) \otimes f_1(C) \\ \downarrow \eta_B & & \downarrow \eta_A \otimes \eta_C \\ f'_1(B) & \xrightarrow{f'_2(\Delta)} & f'_1(A) \otimes f'_1(C) \end{array}$$

is commutative. Before proving this in full generality we first show certain special cases and auxiliary results. To simplify the notation we will write $[A]$ for $f_1(A)$ if A is an object in $\mathcal{T}$ and $[A]$ for $f_1(JA)$ if A is an object in $\text{Gr}^b(\mathcal{E})$. In particular we have $f'_1(A) = f_1(J\tilde{H}A) = [\tilde{H}A]$.

Lemma 5.12. *If $A \in \mathcal{T}^{\geq i} \cap \mathcal{T}^{\leq i}$ and $B, C \in \mathcal{T}^{\geq i}$ then diagram (17) is commutative.*

Proof. We note that the assumptions imply that the boundary homomorphism in the long exact cohomology sequence of Δ (and of all auxiliary distinguished triangles in this proof) is zero.

Case (i): If $A = \tau_{\leq i}B$, $C = \tau_{>i}B$ then the statement follows easily from the definition of η_B .

Case (ii): If $B, C \in \mathcal{T}^{\leq i}$ then the distinguished triangle $A \rightarrow B \rightarrow C \rightarrow TA$ can be identified with the distinguished triangle induced from the short exact sequence $0 \rightarrow \tilde{H}A \rightarrow \tilde{H}B \rightarrow \tilde{H}C \rightarrow 0$, which implies the result.

General case: There is an octahedral diagram

$$\begin{array}{ccccccc}
 A & \longrightarrow & \tau_{\leq i} B & \longrightarrow & \tau_{\leq i} C & \longrightarrow & TA \\
 \parallel & & \downarrow & & \downarrow & & \parallel \\
 A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & TA \\
 & & \downarrow & & \downarrow & & \\
 & & \tau_{> i} B & \xlongequal{\quad} & \tau_{> i} B & & \\
 & & \downarrow & & \downarrow & & \\
 & & T\tau_{\leq i} B & \longrightarrow & T\tau_{\leq i} C & &
 \end{array}$$

and a corresponding exact diagram of cohomology objects. It follows that the front face and back face in the following diagram are commutative.

$$\begin{array}{ccccc}
 & & [\tilde{H}B] & \xrightarrow{f'_2(\Delta)} & [\tilde{H}A] \otimes [\tilde{H}C] \\
 & \nearrow \eta_B & \downarrow & \nearrow \eta_A \otimes \eta_C & \downarrow \\
 [B] & \xrightarrow{f_2(\Delta)} & [A] \otimes [C] & & \\
 \downarrow & & \downarrow & & \downarrow \\
 & \nearrow \eta_{\tau_{\leq i} B} \otimes \eta_{\tau_{> i} B} & [\tilde{H}\tau_{\leq i} B] \otimes [\tilde{H}\tau_{> i} B] & \longrightarrow & [\tilde{H}A] \otimes [\tilde{H}\tau_{\leq i} C] \otimes [\tilde{H}\tau_{> i} B] \\
 & \downarrow & \downarrow & \nearrow \eta_A \otimes \eta_{\tau_{\leq i} C} \otimes \eta_{\tau_{> i} B} & \\
 [\tau_{\leq i} B] \otimes [\tau_{> i} B] & \longrightarrow & [A] \otimes [\tau_{\leq i} C] \otimes [\tau_{> i} B] & &
 \end{array}$$

The left and right hand faces are commutative by case (i) considered above (note that $\tau_{> i} B \cong \tau_{> i} C$ canonically), and the bottom face is commutative by case (ii). It follows that the top face is commutative which is the statement of the lemma. $\square$

For any object A in $\mathcal{T}$ we let μ_A denote the unit structure on $[A] \otimes [TA]$ which is induced by the determinant functor f (compare Lemma 3.6). Using the canonical isomorphism $TJ\tilde{H}A \cong JT\tilde{H}A$, we will consider $\mu_{J\tilde{H}A}$ also as a unit structure on $[J\tilde{H}A] \otimes [JT\tilde{H}A] = [\tilde{H}A] \otimes [T\tilde{H}A]$.

Lemma 5.13. *Let A be an object in $\mathcal{T}$. Then the composite map $[A] \otimes [TA] \xrightarrow{\eta_A \otimes \eta_{TA}} [J\tilde{H}A] \otimes [J\tilde{H}TA] \xrightarrow{\cong} [J\tilde{H}A] \otimes [JT\tilde{H}A]$ (where we use the canonical isomorphism $J\tilde{H}TA \cong JT\tilde{H}A$) is an isomorphism of units $([A] \otimes [TA], \mu_A) \rightarrow ([\tilde{H}A] \otimes [T\tilde{H}A], \mu_{J\tilde{H}A})$.*

Proof. If $A \in \mathcal{T}^{\geq i} \cap \mathcal{T}^{\leq i}$ then the statement is easy to see. In general we use induction on the number of non-zero cohomology objects of A , where the inductive step uses Lemmas 5.12 and 3.6(ii) applied to the distinguished triangles $\tau_{\leq i} A \xrightarrow{a} A \xrightarrow{b} \tau_{> i} A \xrightarrow{c} T\tau_{\leq i} A$ and $T\tau_{\leq i} A \xrightarrow{Ta} TA \xrightarrow{Tb} T\tau_{> i} A \xrightarrow{Tc} T^2\tau_{\leq i} A$. $\square$

Lemma 5.14. *Suppose that the diagram of the form (17) associated to the rotated triangle $\Delta' : B \xrightarrow{b} C \xrightarrow{c} TA \xrightarrow{-Ta} TB$ is commutative. Then the diagram (17) associated to the original triangle $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$ is commutative.*

Proof. Consider the following diagram.

$$\begin{array}{ccccc}
[B] & \xrightarrow{\mu_A} & [A] \otimes [TA] & \xrightarrow{f_2(\Delta')^{-1}} & [A] \otimes [C] \\
\downarrow \eta_B & & \downarrow \eta_A \otimes \eta_{TA} \otimes \eta_B & & \downarrow \eta_A \otimes \eta_C \\
[\tilde{H}B] & \xrightarrow{\mu_{J\tilde{H}A}} & [\tilde{H}A] \otimes [T\tilde{H}A] & \xrightarrow{\alpha_1} & [\tilde{H}A] \otimes [T \ker \tilde{H}a] \otimes [T \ker \tilde{H}b] \\
& & \otimes [\tilde{H}B] & & \otimes [\ker \tilde{H}b] \otimes [\ker \tilde{H}c] \\
& & \downarrow \beta_1 & & \downarrow \beta_2 \\
& & [\ker \tilde{H}a] \otimes [T \ker \tilde{H}a] & \xrightarrow{\alpha_3} & [\ker \tilde{H}a] \otimes [T \ker \tilde{H}a] \\
& & \otimes [\ker \tilde{H}b] \otimes [T \ker \tilde{H}b] & & \otimes [\ker \tilde{H}b] \otimes [T \ker \tilde{H}b] \\
& & \otimes [\tilde{H}B] & & \otimes [\ker \tilde{H}b] \otimes [\ker \tilde{H}c] \\
& & \downarrow \mu_{J \ker \tilde{H}b}^{-1} & & \downarrow \mu_{J \ker \tilde{H}b, \text{top}}^{-1} \\
[\tilde{H}B] & \xrightarrow{\mu_{J \ker \tilde{H}a}} & [\ker \tilde{H}a] \otimes [T \ker \tilde{H}a] & \xrightarrow{\alpha_4} & [\ker \tilde{H}a] \otimes [T \ker \tilde{H}a] \\
& & \otimes [\tilde{H}B] & & \otimes [\ker \tilde{H}b] \otimes [\ker \tilde{H}c] \\
& & & & \xrightarrow{\alpha_5} [\tilde{H}A] \otimes [\tilde{H}C]
\end{array}$$

We will say that a map in the diagram is induced by a short exact sequence Σ in $\text{Gr}^b(\mathcal{E})$ if it is obtained by applying either $f_2(J(\Sigma))$ or $f_2(J(\Sigma))^{-1}$.

The map α_1 is induced by the short exact sequences $\ker \tilde{H}b \rightarrow \tilde{H}B \xrightarrow{\tilde{H}b} \ker \tilde{H}c$ and $T \ker \tilde{H}a \rightarrow T\tilde{H}A \xrightarrow{-T\tilde{H}a} T \ker \tilde{H}b$, and the map α_2 is induced by $\mu_{J \ker \tilde{H}b}^{-1}$ and the short exact sequence $\ker \tilde{H}c \rightarrow \tilde{H}C \xrightarrow{\tilde{H}c} T \ker \tilde{H}a$. The maps α_3 and α_4 are induced by the short exact sequence $\ker \tilde{H}b \rightarrow \tilde{H}B \xrightarrow{\tilde{H}b} \ker \tilde{H}c$, and the map α_5 is induced by the short exact sequences $\ker \tilde{H}a \rightarrow \tilde{H}A \xrightarrow{\tilde{H}a} \ker \tilde{H}b$ and $\ker \tilde{H}c \rightarrow \tilde{H}C \xrightarrow{\tilde{H}c} T \ker \tilde{H}a$.

The map β_1 is induced by the short exact sequences $\ker \tilde{H}a \rightarrow \tilde{H}A \xrightarrow{\tilde{H}a} \ker \tilde{H}b$ and $T \ker \tilde{H}a \rightarrow T\tilde{H}A \xrightarrow{T\tilde{H}a} T \ker \tilde{H}b$. Finally β_2 is induced by $\ker \tilde{H}a \rightarrow \tilde{H}A \xrightarrow{\tilde{H}a} \ker \tilde{H}b$ and $\varepsilon([\ker \tilde{H}b])$.

The top left hand square is commutative by Lemma 5.13, and the top right hand rectangle by our assumption that the statement of the theorem holds for Δ' . The bottom left hand rectangle is commutative because μ is compatible with distinguished triangles, cf. Lemma 3.6(ii). The two squares in the middle are obviously commutative (for the upper one note that the minus sign in α_1 and the $\varepsilon([\ker \tilde{H}b])$ in β_2 cancel). Finally the bottom right hand rectangle is commutative because we are applying $\mu_{J \ker \tilde{H}b}$ to two different copies of $[\ker \tilde{H}b]$ which cancels the $\varepsilon([\ker \tilde{H}b])$ in β_2 .

The composite of the two maps in the top row is $f_2(\Delta)$ (by Lemma 3.8) and the composite of the three maps in the bottom row is the map $f'_2(\Delta) : [\tilde{H}B] \rightarrow [\tilde{H}A] \otimes [\tilde{H}C]$ induced by the long exact cohomology sequence. Hence the commutativity of the outer square is the required result. $\square$

We can now show that diagram (17) is commutative for every distinguished triangle $\Delta : A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$. We do this by induction on the length of the long exact cohomology sequence of Δ , where the length is defined as follows: if we write

the cohomology sequence in the form $\cdots \rightarrow E^{-1} \rightarrow E^0 \rightarrow E^1 \rightarrow E^2 \rightarrow \cdots$ where $E^{-1} = H^{-1}(C)$, $E^0 = H^0(A)$, $E^1 = H^0(B)$, $E^2 = H^0(C)$ etc., then we define the length to be 0 if all E^i are zero objects, and to be $j - i$ if i (resp. j) is the smallest (resp. largest) index for which E^i (resp. E^j) is non-zero.

If the cohomology sequence of Δ has length 0 then the commutativity of (17) is a special case of Lemma 5.12.

Now assume that the cohomology sequence of Δ contains a non-zero object. Lemma 5.14 shows that we can assume that the lowest non-zero term in the cohomology sequence is $H^i(A)$ for some i . We define an object D of $\mathcal{T}$ by the octahedral diagram

$$(18) \quad \begin{array}{ccccccc} \tau_{\leq i} A & \longrightarrow & A & \longrightarrow & \tau_{> i} A & \longrightarrow & T\tau_{\leq i} A \\ \parallel & & \downarrow a & & \downarrow d & & \parallel \\ \tau_{\leq i} A & \longrightarrow & B & \longrightarrow & D & \longrightarrow & T\tau_{\leq i} A \\ & & \downarrow & & \downarrow & & \\ & & C & \xlongequal{\quad} & C & & \\ & & \downarrow & & \downarrow & & \\ & & TA & \longrightarrow & T\tau_{> i} A & & \end{array}$$

where the first vertical triangle is Δ and the first horizontal triangle is the canonical one. We denote the second vertical triangle by Δ' and note that the cohomology sequence of Δ' is shorter than the cohomology sequence of Δ . From (18) we obtain the following commutative exact diagram in $\text{Gr}^b(\mathcal{E})$.

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \ker \tilde{H}a & \xlongequal{\quad} & \ker \tilde{H}d & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T^{-i}H^i A & \longrightarrow & \tilde{H}A & \longrightarrow & \tilde{H}\tau_{> i} A \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T^{-i}H^i A & \longrightarrow & \tilde{H}B & \longrightarrow & \tilde{H}D \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & \tilde{H}C & \xlongequal{\quad} & \tilde{H}C \\ & & & & \downarrow & & \downarrow \\ & & & & T \ker \tilde{H}a & \xlongequal{\quad} & T \ker \tilde{H}d \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

Finally consider the following diagram in $\mathcal{P}$.

$$\begin{array}{ccccc}
& & [\tilde{H}B] & \xrightarrow{f'_2(\Delta)} & [\tilde{H}A] \otimes [\tilde{H}C] \\
& \nearrow \eta_B & \downarrow & & \nearrow \eta_A \otimes \eta_C \\
[B] & \xrightarrow{f_2(\Delta)} & [A] \otimes [C] & & \\
\downarrow & & \downarrow & & \downarrow \\
& & [T^{-i}H^i A] \otimes [\tilde{H}D] & \xrightarrow{\text{id} \otimes f'_2(\Delta')} & [T^{-i}H^i A] \otimes [\tilde{H}\tau_{>i}A] \otimes [\tilde{H}C] \\
& \nearrow \text{id} \otimes \eta_D & \downarrow & & \nearrow \text{id} \otimes \eta_{\tau_{>i}A} \otimes \eta_C \\
[\tau_{\leq i}A] \otimes [D] & \xrightarrow{\text{id} \otimes f_2(\Delta')} & [\tau_{\leq i}A] \otimes [\tau_{>i}A] \otimes [C] & &
\end{array}$$

The commutativity of the bottom face follows from the induction hypothesis applied to $\Delta' : \tau_{>i}A \rightarrow D \rightarrow C \rightarrow T\tau_{>i}A$, and the commutativity of the back face follows easily from the exact commutative diagram above. The front is commutative by the associativity axiom applied to diagram (18). The commutativity of the left and right hand faces are special cases of Lemma 5.12. The resulting commutativity of the top face gives the commutativity of diagram (17) and therefore completes the proof of Proposition 5.11. $\square$

REFERENCES

- [1] A.-A. Beilinson, J. Bernstein, P. Deligne, *Analyse et topologie sur les espaces singuliers I.*, Astérisque **100** (1982).
- [2] M. Breuning, D. Burns, *Additivity of Euler characteristics in relative algebraic K-groups*, Homology, Homotopy and Applications, **7** no. 3 (2005), 11–36.
- [3] D. Burns, *Equivariant Whitehead torsion and refined Euler characteristics*, Number theory, 35–59, CRM Proc. Lecture Notes, **36**, Amer. Math. Soc., Providence, RI, 2004.
- [4] D. Burns, M. Flach, *Tamagawa numbers for motives with (non-commutative) coefficients*, Doc. Math. **6** (2001), 501–570.
- [5] P. Deligne, *Le déterminant de la cohomologie*, Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), 93–177, Contemp. Math. **67**, Amer. Math. Soc., Providence, RI, 1987.
- [6] T. Fukaya, K. Kato, *A formulation of conjectures on p-adic zeta functions in non-commutative Iwasawa theory*, to appear in Proc. St. Petersburg Math. Soc. **11** (2005), english translation to be published by the Amer. Math. Soc.
- [7] F. F. Knudsen, *Determinant functors on exact categories and their extensions to categories of bounded complexes*, Michigan Math. J. **50** (2002), no. 2, 407–444.
- [8] F. F. Knudsen, D. Mumford, *The projectivity of the moduli space of stable curves. I: Preliminaries on "det" and "Div"*, Math. Scand. **39** (1976), no. 1, 19–55.
- [9] F. Muro, A. Tonks, *The 1-type of a Waldhausen K-theory spectrum*, preprint 2006, [arXiv:math.KT/0603544](https://arxiv.org/abs/math.KT/0603544).
- [10] A. Neeman, *The K-theory of triangulated categories*, in E. M. Friedlander, D. R. Grayson (eds.), *Handbook of K-theory*, Springer-Verlag, 2005.
- [11] N. Saavedra Rivano, *Catégories Tannakiennes*, Lecture Notes in Mathematics **265**, Springer-Verlag, 1972.
- [12] J. L. Verdier, *Catégories dérivées, état 0*, in *SGA 4 1/2*, Lecture Notes in Mathematics **569**, Springer-Verlag, 1977.

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, STRAND, LONDON WC2R 2LS,
UNITED KINGDOM

E-mail address: manuel.breuning@kcl.ac.uk